



# The photosynthesis game is in the "inter-play": Mechanisms underlying CO<sub>2</sub> diffusion in leaves



J. Gago<sup>a</sup>, D.M. Daloso<sup>b</sup>, M. Carriqui<sup>a,c</sup>, M. Nadal<sup>a</sup>, M. Morales<sup>a</sup>, W.L. Araújo<sup>d</sup>, A. Nunes-Nesi<sup>d</sup>, A.V. Perera-Castro<sup>a</sup>, M.J. Clemente-Moreno<sup>a</sup>, J. Flexas<sup>a,\*</sup>

<sup>a</sup> Research Group on Plant Biology Under Mediterranean Conditions, Departament De Biologia, Universitat de les Illes Balears / Institute of Agro-Environmental Research and Water Economy –INAGEA, Carretera de Valldemossa, 07122, Palma de Mallorca, Spain

<sup>b</sup> Departamento de Bioquímica e Biologia Molecular, Universidade Federal do Ceará, Fortaleza-CE, 60451-970, Brazil

<sup>c</sup> School of Natural Sciences, University of Tasmania, Bag 55, 7001, Hobart, Tasmania, Australia

<sup>d</sup> Departamento de Biologia Vegetal, Universidade Federal de Viçosa, 36570-900, Viçosa, Minas Gerais, Brazil

## ARTICLE INFO

### Keywords:

Photosynthesis  
Water use efficiency  
Stomatal conductance  
Mesophyll conductance  
Diffusional limitations

## ABSTRACT

Ensuring global food security is a worldwide major concern considering the predicted climate change scenarios for the main agricultural regions of the world. Stomatal conductance ( $g_s$ ) and mesophyll conductance ( $g_m$ ) are major drivers limiting photosynthesis (A). Both conductances frequently impose about two-thirds of the total photosynthetic limitation under optimum conditions. However, under abiotic stress, like drought or salinity, the diffusional limitations can reach more than 85 % of the total. Thus, knowledge about both conductances is essential to improve water use efficiency (WUE) through targeted crop breeding programs and to promote sustainable, resource-efficient, and environmental-friendly agriculture strategies.

Intriguingly, knowledge obtained from both conductances from decades of research differs importantly by their focus. Whilst the role of both mechanics and metabolism of guard cells on the regulation of stomatal movements remains much less understood, the signaling pathways that regulates stomatal movements are well-documented. The opposite is true for  $g_m$ , in which the biochemical regulation and signaling pathways remain mostly still unexplored. Even more surprising is the lack of information about the putative molecular mechanisms that should drive the known coupled behavior of both conductances in response to the environment. Here, we discussed the main mechanisms driving the responses of each of the conductances, but highlighting a special focus into the possible common determinants that could link their coupled behavior. Further integrative multidisciplinary studies joining molecular biology and ecophysiology are required in order to improve our understanding of both conductances, the major actors limiting photosynthesis and WUE in a changing environment.

## 1. Out of the atmosphere and into the leaf: the complex CO<sub>2</sub> pathway to photosynthesis

The origin and early diversification of land plants began in the mid-Paleozoic era (ca. 470 Ma), where earlier plants faced new conditions in an aerial environment, requiring adaptations to obtain a positive carbon balance without being dehydrated out of the aquatic environment (Kenrick and Crane, 1997; Quirk et al., 2014). About 20 million years later, flat-bladed megaphylls, highly specialized photosynthetic structures resembling the so-called 'leaves' (*sensu lato*), suddenly spread throughout land floras (Osborne et al., 2004). Another unprecedented evolutionary innovation was the development of impermeable cuticles, closely followed by the appearance of the stomata ca. 400 Ma,

consisting of the stomatal pore surrounded by two guard cells mainly at the leaf epidermis. While it appears that, originally, the main function of stomata was to regulate water loss allowing, for instance, sporophytes to desiccate (Duckett and Pressel, 2018; Brodribb et al., 2020), this exclusive evolutionary innovation was soon also used by leaves to regulate the CO<sub>2</sub> and water exchange between the leaf and the surrounding atmosphere (Edwards et al., 1998). It is still a matter of debate how leaf evolution, stomata, and the changing climatic conditions interacted and influenced each other leading to the current leaf structure types and functions (Osborne et al., 2004; Franks and Beerling, 2009; Leakey and Lau, 2012). Nevertheless, compelling evidence suggests that stomata have appeared before bryophytes, although the true identity of the stomata ancestor remains unknown (Harris et al., 2020).

\* Corresponding author.

E-mail address: [jflexas@uib.es](mailto:jflexas@uib.es) (J. Flexas).

<https://doi.org/10.1016/j.envexpbot.2020.104174>

Received 22 February 2020; Received in revised form 24 June 2020; Accepted 1 July 2020

Available online 02 July 2020

0098-8472/© 2020 Published by Elsevier B.V.

Stomata most likely represent the very first step for the photosynthetic  $\text{CO}_2$  pathway in vascular plants, and their highly regulated opening / closing leads to a largely variable and dynamic stomatal conductance ( $g_s$ ). Subsequently, the  $\text{CO}_2$  reaches the substomatal cavities at a certain concentration ( $C_i$ ). Openness of the stomata pore has additional consequences including transpiration of the water vapour present in sub-stomatal cavities ( $W_i$ ). Thus, the balance among leaf carbon gain and water loss constitutes an essential relationship for plants living in environments with limited water, such as those with important abiotic stresses like drought, salinity and/or high atmospheric vapor pressure deficit. This relationship is known as the water use efficiency (WUE), an essential parameter for agriculture that can be considered in different spatial and temporal scales, and becoming even more important now considering the climatic change predicted scenarios for the agricultural regions (Gago et al., 2014; WWAP, 2018; Leakey et al., 2019).

From the substomatal cavities, the  $\text{CO}_2$  diffuses into the mesophyll tissues to the carboxylation sites in the chloroplast, the so-called mesophyll conductance ( $g_m$ ). This pathway is highly complex because  $\text{CO}_2$  should cross different media within the leaves such as airspaces, water and biophysical barriers (such as cell walls -CWs- and membranes), and, moreover, it can be actively regulated by e.g. transmembrane channels like aquaporins and enzymes like carbonic anhydrases to finally reach the Rubisco in the stroma (Flexas et al., 2012). The gas-phase within the mesophyll is known as the intercellular air spaces diffusion conductance ( $g_{ias}$ ), that is driven by the effective mesophyll thickness and porosity to the outer surface of the CWs. Subsequently, the liquid phase consists of a series of conductances (i.e. the inverse of resistances) including the cell wall ( $g_{cw}$ ), the plasma membrane ( $g_{pm}$ ), the cytoplasm ( $g_{cyt}$ ), the chloroplast envelope ( $g_{env}$ ) and the chloroplast stroma ( $g_{st}$ ) itself (Terashima et al., 2011), where finally  $\text{CO}_2$  reaches the carboxylation sites in the stroma at a certain concentration ( $C_c$ ).

While  $g_s$  has been measured repeatedly since the onset of gas exchange systems (Gaastra, 1959; Björkman and Holmgren, 1963), it is only more recently that estimates of  $g_m$  have been obtained, combining gas exchange measurements with either on-line carbon isotope discrimination (Evans et al., 1986) or chlorophyll fluorescence (Harley et al., 1992) (see Pons et al., 2009 for a review on currently used techniques).

Leaf photosynthesis ( $A$ ) is totally dependent on the  $\text{CO}_2$  availability in the sites of carboxylation and it is thus strongly influenced by both  $g_s$  and  $g_m$ . Hence, high photosynthetic capacity is expected to be associated to high  $g_s$  and  $g_m$  (Flexas et al., 2013a,2013b; Onoda et al., 2017;


Nadal and Flexas, 2019), and a positive relationship between  $g_s$  and  $g_m$  can be found across species under optimum conditions (Flexas et al., 2013a; Nadal and Flexas, 2018). Nonetheless, there is still a significant variation in the  $g_m/g_s$  ratio across species and even genotypes (Flexas et al., 2013a,2013b), and this ratio changes in responses to abiotic factors like water availability (Flexas et al., 2013a) or plant's nutritional status (Franco-Navarro et al., 2019). This variation has to be considered an important feature for e.g. selection of genotypes based on WUE, especially under drought conditions (Galmés et al., 2011; Thérour-Rancourt et al., 2015). Furthermore, the dynamic behavior of both conductances plays a major role in driving terrestrial primary productivity, but it is still a great challenge to implement the leaf-scale knowledge concerning both conductances into ecosystems and global models. The challenge for modelling studies will be thus to integrate the dynamic of  $g_s$  and  $g_m$  responses with different remote sensing technologies in order to improve both carbon and water flux predictions at a global scale (Guanter et al., 2014; Sun and Gu, 2014; Knauer et al., 2019, 2020).

In this review, we will focus in the latest advances about the mechanisms driving both conductances ( $g_s$  and  $g_m$ ) as the major drivers of the  $\text{CO}_2$  availability at the carboxylation site and leaf-level WUE. In addition, we discuss the essential metabolism interplay between them in an attempt to improve the current understanding of the complex physiological responses of both conductances and photosynthesis.

## 2. The importance of $g_s$ and $g_m$ and current knowledge on their regulation

As outlined in the above section,  $g_s$  and  $g_m$  similarly limit photosynthesis in higher plants. However, as recently shown by Gago et al. (2019), their importance as photosynthesis-limiting factors is variable across the land plant's phylogeny in the absence of stress conditions and at saturating light. Combined,  $g_s$  and  $g_m$  account for more than 75 % of the total limitation in bryophytes, lycophytes, and pteridophytes, and they account for about 70 % in gymnosperms. In angiosperms, on average, each of the three limiting factors account for about 1/3 of the total limitation, that is, combined diffusional limitations account for 2/3 while photo-biochemical limitations account for only 1/3 of the total limitation. Under abiotic stress e.g. water or salt stress, the diffusional limitations typically increase far more than photo-biochemical limitations, hence accounting for an even larger fraction of total limitations (Flexas et al., 2006; Chaves et al., 2009).

The well-proven importance of the two diffusional photosynthetic



**Fig. 1.** A summary of current knowledge on the regulation of both stomatal and mesophyll conductance. The size of boxes and letters is proportional to the magnitude of current knowledge on each case. Hence, for stomatal conductance its structural determinants are largely unknown but there is abundant knowledge both on signalling pathways and biochemical regulation. The opposite is true for mesophyll conductance, for which there is large knowledge on its structural determinants and very little is known about both signalling and biochemical regulation. Abbreviations: ABA, abscisic acid; IAS, intercellular air spaces, SLAC1, Slow Anion Channel Associated 1 (in the plasma membrane); PP2Cs/SnRK2, phosphatases/kinases regulation (in response to ABA in guard cells).

limitations contrasts with the scarce knowledge available on some aspects of their respective regulation (Fig. 1). Although the importance of stomata morpho-anatomical features, such as size and distribution, on the regulation of stomatal conductance have been widely investigated (Woelfenden et al., 2018; Zoulias et al., 2018), knowledge concerning the signaling pathways that regulate stomatal movements are also significantly understood. However, whereas the major structural determinants of  $g_m$  are well-studied, the molecular and biochemical mechanisms behind  $g_m$  regulation remain rather unexplored. In the following sections, we will summarize the current knowledge of the anatomical and molecular determinants of  $g_s$  (Section 3) and  $g_m$  (Section 4) including their responses to the environment, and finally the coupled behavior and interplay between both conductances driving photosynthesis (Section 5).

### 3. Stomatal conductance: the leaf gates for photosynthesis

Stomata is a key structure for plant environmental signal perception. Environmental cues such as light,  $CO_2$  concentration and water availability are known to impact both stomatal movement and density (Casson and Hetherington, 2010). These signals are directly perceived by guard cells and indirectly through signalling molecules that tightly connect mesophyll photosynthetic activity with guard cells function. At geological time-scale, it is believed that the  $CO_2$  atmospheric concentration was a major force driven stomata morphology and physiology (Franks and Beerling, 2009; Brodribb and McAdam, 2017). For instance, the decrease in environmental  $CO_2$  concentration leads to decreased stomatal size and increased stomatal density, which is believed to be a response to optimize the flux of  $CO_2$  from the atmosphere to the internal leaf space (Franks and Beerling, 2009). Additionally, it has been shown that stomatal movements have acquired an active control over plant phylogenetic groups, resulting in angiosperms with faster stomatal responses, in which the main stress-related phytohormone abscisic acid (ABA) play a central role in coordinating stomatal responses to closure stimulus such as drought, darkness or high  $CO_2$  concentration (Assmann and Jegla, 2016). Deciphering stomata evolution and functionality promoted a great interest from molecular to ecophysiological perspectives in the last decades (Doi et al., 2006; Doi and Shimazaki, 2008; McAdam and Brodribb, 2012a, 2012b; Franks and Britton-Harper, 2016; Deans et al., 2019). Whilst some authors have suggested a gradualistic evolution of stomatal control (Brodribb and McAdam, 2011), others suggest that ancient stomata from basal land plant groups already possess all mechanisms regarding sensitivity and response to the environment as the modern angiosperms (Ruszcza et al., 2011; Franks and Britton-Harper, 2016; Chater et al., 2017). In any case, despite some common mechanisms of stomatal regulation, tremendous differences in both stomatal behaviour and morphology are found among the different plant groups. It is clear that basal lineage of plants (e.g. ferns) have much lower  $g_s$ , which is, at least to some extent, associated with a lower stomatal density (Lima et al., 2019). However, it is noteworthy that significant differences in stomata morphology are also found even within angiosperm species (McAusland et al., 2016; Lawson and Viallet-Chabrand, 2018). For instance, whilst eudicots typically present kidney-shaped stomata, monocots can have either kidney-shaped or dumbbell-shaped stomata with adjacent subsidiary cells (Bertolino et al., 2019). Therefore, it seems that the combination of patterning of stomata formation and regulation follows specific environmental requirements toward plant WUE improvement.

Several recent evidence indicate that stomata from ferns are able to respond to changes in light/dark and different  $CO_2$  concentrations, although at a much lower velocity (Franks and Britton-Harper, 2016; Lima et al., 2019). Thus, the main achievement of angiosperms stomata is the ability to rapidly respond to environmental cues (McAusland et al., 2016; Lawson and Viallet-Chabrand, 2018). Faster stomata in angiosperms could be associated to genetic modifications such as the appearance of new genes or simply the duplication of a function of old

ones coupled to anatomical and physiological differences among these and basal plants (Lima et al., 2019; Susmilch et al., 2019a, b). Given that the velocity of stomatal responses are pivotal for plant responses to abiotic stress and improving WUE (Lawson and Blatt, 2014; Lawson and Viallet-Chabrand, 2018), future studies aiming to unveil the evolution and the regulatory mechanisms of stomatal speediness are of crucial importance to improve plant WUE, particularly in crops. Whilst slower stomatal opening may lead to unnecessary water losses and improved WUE in a scenario in which  $g_s$  is partially uncoupled to  $A$  (McAusland et al., 2016), plants with faster stomatal closure would avoid water loss during stress conditions. Compelling evidence has recently revealed that guard cell starch degradation is a key regulator of light stomatal speediness (Flütsch et al., 2020). It remains unclear, however, how and to which extent this mechanism affects plant performance under fluctuating environmental conditions.

#### 3.1. The determinants of maximum stomatal conductance

Under optimal conditions (i.e. well-hydrated plants, saturating light and low vapour pressure deficits), leaves are able to attain a maximum stomatal conductance ( $g_{smax}$ ). Potential  $g_{smax}$  is achieved by anatomically-driven traits which are established during leaf development: the stomatal density ( $D_s$ , i.e. the number of stomata per unit leaf surface) and the stomatal pore depth and maximum area, which are related to size ( $S_s$ ) and the shape of stomata (Franks and Beerling, 2009; Sack and Buckley, 2016; Harrison et al., 2019). Higher  $g_{smax}$  values are achieved by increases in  $D_s$ , which are often correlated with decreases in  $S_s$ , coordination that has been widely observed as a result of both acclimation and adaptation responses to variations in environmental factors (Franks and Beerling, 2009; Doherty-Adams et al., 2012; Harrison et al., 2019). Coordination between  $D_s$  and  $S_s$  is apparently under tight selection in angiosperms, determining the optimal use of the epidermal area for  $g_{smax}$  (de Boer et al., 2016a, 2016b; Haworth et al., 2018). Only a few species, like some species within *Begonia* genus, have been able to overcome the minimum one-epidermal cell spacing between stomata (Gan et al., 2010; Harrison et al., 2019). Regarding stomatal shapes, which are species-specific but diverse across land plants, some allow opening comparatively wider than others hence being able to achieve higher  $g_{smax}$  values. For instance, in angiosperms, dumbbell-shaped stomata allow for wider opening than kidney-like-shaped stomata; and kidney-shaped stomata with thinner guard cells allow for wider opening than the same with wider guard cells (Franks, 2006; Franks and Farquhar, 2007).

Leaves do not usually operate at  $g_{smax}$  and thus  $g_s$  typically remains at around 20–30 % of  $g_{smax}$ , reaching for some angiosperm species up to 39–43 %, corresponding to a turgor pressure at which guard cells more efficiently control the aperture of the pore (Dow and Bergmann, 2014; McElwain et al., 2016). Even so, it sets a maximum evaporative demand that needs to be matched by a hydraulic supply of the same magnitude. Otherwise, this would result in damage to the water transport tissues, potentially even causing the embolism of the vascular tissue (Brodribb and Feild, 2000; Hubbard et al., 2001; Brodribb and Holbrook, 2004; Brodribb and Buckley, 2018). Thereby, as the leaf hydraulic capacity should coordinate with  $g_{smax}$ , the optimization to achieve higher  $CO_2$  diffusion rates of the stomatal properties and density came necessarily associated with a concomitant increase of this hydraulic supply over the more of 400 Myr of land plant evolution (Brodribb et al., 2020). This coordination occurs, and there is a conservative investment in the water conductive tissues to fair enough supply  $g_{smax}$  (Medlyn et al., 2011; Martins et al., 2014; Carins Murphy et al., 2016; Brodribb and McAdam, 2017). Thus, since leaves impose a disproportionately large resistance to the hydraulic system (Sack and Holbrook, 2006), leaf hydraulic conductance ( $K_{leaf}$ ) is usually tightly correlated to  $g_s$  and photosynthesis (Brodribb et al., 2007; Scoffoni et al., 2016). From a mechanistic perspective, the leaf hydraulics link with  $g_{smax}$  and  $D_s$  is phylogenetically mostly driven by a positive correlation with vein density ( $D_v$ ) (Boyce

et al., 2009; McElwain et al., 2016), although they do not always correlate at the short phylogenetic scale (Gleason et al., 2019; McElwain et al., 2016; Rockwell and Holbrook, 2017).

In summary,  $g_{smax}$  is mainly anatomically determined by a link between  $D_s$  and  $D_v$  and through changes in the leaf cell size (i.e. miniaturizing the size of stomata and veins, but also of other leaf cell types) (Brodribb and McAdam, 2013; Feild and Brodribb, 2013; Carins Murphy et al., 2016; de Boer et al., 2016a, 2016b; Carins Murphy et al., 2017). Interestingly, this, in turn, appears to be scaled by the genome size (Simonin and Roddy, 2018; Roddy et al., 2018). However, despite the great advance in our understanding concerning stomatal development and evolution, how the different stomata morphologies associated with their intrinsic hydraulic and biochemical regulation contribute to stress tolerance remains far from clear (Medeiros et al., 2019). This is particularly due to the fact that the regulation of stomatal movements from highly stress-tolerant groups such as bryophytes, ferns, resurrection and CAM plants remains poorly and fragmentarily studied (Males and Griffiths, 2017), which clearly hampers our understanding on how stomatal physiology could be used in plant breeding programs toward plant stress tolerance.

### 3.2. Stomatal responses to the environment: perception and signalling

Changes in guard cell metabolism are known to regulate the magnitude of stomatal opening according to the prevailing environmental conditions (Lawson et al., 2014a). Guard cells are thus able to directly recognize light,  $CO_2$  and other signals and then modulate stomatal opening or closure. In general, stomata open in the light and close during dark conditions, with the exception of CAM plants. However, a significant transpiratory stream is detected in the night for many species independent of their origin, and the precise function of this nocturnal conductance ( $g_n$ ) is still a matter of debate: why plants sustain  $g_n$  under dark conditions losing water without carbon gains? Recently, through a multi-species meta-analysis, it was demonstrated that  $g_n$  was related to the relative growth rate (RGR) (Resco de Dios et al., 2019). These results do not relate with the previous hypothesis explaining the  $g_n$  role as the removal of  $CO_2$  excess produced by respiration,  $O_2$  delivery to the vascular system, nutrient supply or stomata leakiness. Notably, it has been postulated that higher  $g_n$  would optimize predawn  $g_s$  and subsequently earlier carbon gains (Resco de Dios et al., 2019 and references therein). Curiously, it appears that  $g_n$  responds roughly to the very same environmental factors as day-time  $g_s$  (Caird et al., 2007). Nevertheless, and besides the still not fully understood role of  $g_n$ , most of our current knowledge on stomatal responses to environmental cues refer to day-time dynamics.

The dynamics of stomatal movement during the day follow a circadian rhythm with a further complex network of interaction with endogenous signals, including the synthesis and transport of phytohormones, ions, metabolites and possibly electrical signals and still highly unknown from mesophyll to guard cells (Gallé et al., 2015; Daloso et al., 2017). Guard cells are thus the hub that integrates environmental and endogenous signals to regulate stomatal movements (Mott, 2009). During the diel course, signals from mesophyll cells are key to close the stomata in periods of high A, as a mechanism to optimize plant WUE. Some authors indicated that these molecules could be a vapour-phase ions, possibly hydronium ions able to change the pH of the guard cells (Mott et al., 2014). Although the real identity of these signals still remains to be identified, additional evidence suggests that the compounds linking mesophyll and with guard cells could range in the size of 100–500 Da. This idea is based in the fact that stomatal responses are disrupted when a membrane excluding molecules in this size range are placed between mesophyll and guard cells (Fujita et al., 2019), which may include sucrose (342 Da), malate (134 Da) and ABA (264 Da). In fact, genetic manipulation of sucrose and malate metabolism leads to altered stomatal movements (Laporte et al., 2002; Nunes-Nesi et al., 2007; Araújo et al., 2011; Antunes et al., 2012,

2017a; Kelly et al., 2013; Lugassi et al., 2015; Daloso et al., 2016b; Wang et al., 2019).

Beyond endogenous signals, it is interesting to highlight that environmental cues such as water availability as well as the light quantity and quality strongly influence  $g_s$ . Water deficit is a well-recognized signal that induces stomatal closure via ABA and perhaps hydraulic and electrical signals (see next Sections 3.2.1 and 3.2.2, respectively), being an important mechanism that restricts  $CO_2$  assimilation in periods of moderate stress (Galmés et al., 2007; Flexas et al., 2009a, 2009b). By contrast, light is the main opening signal where red light was for long believed to influence stomatal behaviour via mesophyll photosynthetic activity (Baroli et al., 2007). However, recent evidence indicates that red light can be directly perceived by guard cells and induces plasma membrane  $H^+$ -ATPases phosphorylation (Ando and Kinoshita, 2018), similarly to the well-described blue-light signalling pathway (Inoue and Kinoshita, 2017). Taken all the endogenous and environmental cues that modulate  $g_s$  together, it is clear that  $g_s$  is very dynamic and highly dependent on the prevailing environmental conditions, in which guard cells act as sensors to integrate all hydraulic, metabolic and light signals to modulate  $g_s$  under both non-stress and stress conditions.

#### 3.2.1. Hydraulic signalling

Early works assumed a hydro-passive mechanism for stomatal closure under water stress and increased vapour pressure deficit (VPD), which intrinsically implies interplay between plant hydraulics and transpiration (see Jones, 1998 for a review). In essence, whenever transpiration exceeds the hydraulic capacity of the plant to supply water to leaves, a water deficit occurs in these resulting in decreased leaf water potential and turgor pressure in the guard cells, thus provoking stomatal closure. It has been proposed that such hydro-passive mechanism operates mostly in plants belonging to phylogenetic groups such as lycophytes and pteridophytes (Cardoso et al., 2019), while the ABA-based hydro-active signalling described in the above section is the main system used by seed plants (Brodribb and McAdam, 2011). However, a water potential-based mechanism for the regulation of  $g_s$  in response to irradiance and other factors was also proposed (Pieruschka et al., 2010). Additionally, recent evidence indicates that VPD-stomatal responses appears to involve a passive hydraulic mechanism as well as a signalling pathway that entails OST1 (Open Stomata 1) and SLAC1 (Slow anion channel 1), which would be controlled by either ABA or by a still unknown ABA-independent mechanism (Merilo et al., 2018).

Besides the hydro-passive mechanism, direct evidence for xylem cavitation driving stomatal closure has been provided. For instance, Salleo et al. (2000) showed that dehydrating air-detached leaves increased stomatal aperture (i.e. the so-called ‘wrong-way response’ or ‘Iwanoff effect’) only after the threshold water potential for the onset of petiole cavitation was reached, after what stomata progressively closed thereafter. Moreover, during a transition from darkness to light, attached leaves initially showed progressively increased stomatal opening and decreased water potential, with no cavitation detected by ultrasonic methods. When the water potential for the onset of petiole cavitation was reached  $g_s$  was at its maximum and cavitation events started being recorded ultra-acoustically. Thereafter,  $g_s$  slightly decreased to reach a steady-state value corresponding to a constant water potential just at the onset of petiole cavitation and the cease of cavitation events (Salleo et al., 2000). Hubbard et al. (2001) provided direct evidence for the involvement of cavitation in stomatal closure. After carefully removing bark and phloem from pine stems, they were able to directly inject discrete amounts of air into the xylem to provoke different degrees of cavitation. After each step injection, an immediate decrease of  $g_s$  was recorded, resulting in a perfect linear correlation between  $g_s$  and the hydraulic conductance when pooling data for all the injection steps (Hubbard et al., 2001).

Stomata water relation is a challenging topic where several authors proposed recent models based on mechanisms related to ABA, hydraulic and osmotic potential, among others, but it is still needed much more



knowledge to develop reliable mechanistic models about stomatal responses to water status (reviewed in Buckley, 2019). Regardless of the mechanism, there is compelling evidence for the existence of a hydraulically-based signalling pathway inducing stomatal closure. The predominance of hormonal *versus* hydraulic stomatal regulation in higher plants is currently the subject of ample debate (Wang et al., 2018; Flexas et al., 2018a, 2018b). While the dominance of either mechanism could be species-specific (Flexas et al., 2018a, 2018b), it is more likely that there is an interplay between the two. In grapevines, for instance, Pou et al. (2008) showed that ABA-based  $g_s$  regulation was dominant during the imposition of drought while hydraulic-based regulation was the most important during the subsequent recovery. Additionally, it was shown that grapevines stomatal closure is induced by hydraulic signals but maintained by ABA during drought (Tombesi et al., 2015). Another example of the interplay between the two mechanisms is the indirect involvement of ABA in embolism refilling (Secchi et al., 2013).

### 3.2.2. Electrical signalling

The operation of electrical signaling controlling stomatal aperture has been much less studied than the hormonal and hydraulic ones. Two different types of electrical signals have been described in plants: action potentials and variation potentials, both involved in the regulation of multiple metabolic and developmental effects (see Fromm and Lautner, 2007 and Gallé et al., 2013 for reviews). As it was known for long that watering a dry root is an efficient way of producing action potentials (Pickard, 1973), Fromm and Fei (1998) followed the concomitant variation of  $g_s$  and photosynthesis with action potentials during water stress and re-watering of maize plants. They found a perfect co-variation among these parameters, action potential decreasing in parallel to  $g_s$  during the imposition of water stress and peaking strong – and preceding stomatal opening by about 15 min – immediately upon re-watering. The latter result led them to hypothesize that action potentials induced by re-watering roots were directly responsible for stomata re-opening. To test this hypothesis, Grams et al. (2007) used a design consisting of potted maize plants with the pots inside a root pressure chamber – to compensate / eliminate hydraulic signals – and a part of the leaf cooled by two aluminum blocks that were cooled by flushing with a mixture of glysanthin/water – to eliminate the propagation of action potentials through the leaf. Using this design, and measuring variations of leaf turgor, action potential and  $g_s$ , they observed that upon re-watering stomata first started closing (another ‘wrong-way response’) in a mechanism regulated by hydraulic signals (turgor changes), to then start opening after several minutes in a mechanism regulated by action potentials.

In addition to these examples related to water stress and recovery, Gallé et al. (2013) showed in soybean that wound-induced action potentials in a leaflet provoked a transient stomatal opening in the adjacent leaflets of the leaf and a direct and large decrease of mesophyll conductance and photosynthesis, i.e. a transitory uncoupling of  $g_s$  and  $g_m$ . It seems reasonable to assume that more studies are clearly required to fully disentangle the role of electrical signalling in controlling  $g_s$  (and  $g_m$ ).

### 3.2.3. Hormonal signalling

As several physiological processes, the stomatal movement is also highly controlled by phytohormones, in which especially ABA, have been closely associated with the formation of stomata (Chater et al., 2017). Remarkably, an evident cross-talk among stomata formation and stomatal movement regulation is usually observed, in which the role of ABA, auxin, brassinosteroids, gibberellins, ethylene, cytokinins, and jasmonic acid, as well as the genes associated with stomatal development, have been revealed (Bergmann, 2006; Qi and Torii, 2018). Both the generation and the regulation of stomatal movements are extremely sensitive to changes in the environment around the leaf and, broadly characterized and especially rich in complexity, involving gene

expression, post-translational modifications, protein-protein interaction, synthesis of signalling compounds such as nitric oxide and reactive oxygen species (ROS) as well as ion transporter activity (Kim et al., 2010). In this vein, the role of ABA is well-documented and will be thus the focus of this section.

ABA is the main phytohormone that connects plant water potential with stomatal movement (Buckley, 2019). Guard cells have been used as a model for the establishment of the ABA signalling pathway, in which the complete signalling pathway including PYR/PYL/RCAR receptors and the upstream components have been identified and confirmed to act in tandem in vitro (Fujii et al., 2020; Park et al., 2009; Santiago et al., 2009; González-Guzmán et al., 2012). Given that plants subjected to water deficit conditions have stomatal closure as one of the main physiological responses, which is usually associated with substantial increases in ABA levels, ABA has been associated to be an important regulator especially under stress conditions (Jarzyniak and Jasinski, 2014). Notwithstanding, ABA is seemingly important also under non-stress conditions (Yoshida et al., 2019b), possibly as an important regulator of transpiration, stomatal density (Hetherington and Woodward, 2003; Lake and Woodward, 2008) and the primary metabolism, as recently has been demonstrated (Yoshida et al., 2019c, a).

ABA has also been pointed as an important contributor to the increased velocity of stomatal responses found in angiosperms when compared to ferns. This idea comes from several studies that show a lack or, at maximum, a slower stomatal  $CO_2$  and ABA responses observed in ferns (Brodribb and McAdam, 2011; McAdam and Brodribb, 2012a, 2012b; Franks and Britton-Harper, 2016; McAdam et al., 2016; Hörak et al., 2017; Lima et al., 2019). Although key genes of the ABA signaling pathway have also been identified in ferns, it seems likely that they have acquired their role in the control of stomatal movements only in recent evolutionary groups (McAdam et al., 2016; Hedrich and Geiger, 2017). Thus, it seems that ABA acts as an important modulator of the changes in stomatal density and stomatal responsiveness to environmental cues, especially the changes in  $CO_2$  concentration observed at a geological scale. Notably, it suggests that ABA determines not only the plant  $CO_2$  responses but also strongly influencing the earth's  $CO_2$  biogeochemistry cycle (Hetherington and Woodward, 2003; Lima et al., 2018).

### 3.3. Biochemical regulation of stomata movements

Guard cells have been widely used to understand the molecular and signalling mechanisms of hormonal regulation, especially those associated with ABA metabolism (Assmann and Jegla, 2016). This hormone acts as a hub in the molecular network associated with stomatal closure induced by drought, low humidity, high  $CO_2$  and dark (Kim et al., 2010; Bauer et al., 2013a, 2013b; Engineer et al., 2016b). After its perception by PYR/PYL/RCAR receptors, ABA response is triggered by inhibition of PP2C phosphatases, activation of different kinases such as SnRK2.6/Srke/OST1, synthesis of important second messengers such as ROS, nitric oxide and  $Ca^{+2}$  and nitric oxide and, ultimately, activation of ion channels such as QUAC1, SLAH3 and SLAC1 (Yoshida et al., 2019b). It is noteworthy, however, that each stimulus activates particular signaling pathways culminating in stomatal closure.

Beyond that, evidences suggest that stomatal closure is regulated by an intricate regulatory network that involves not only ABA but also ethylene, brassinosteroid, jasmonic acid among other phytohormones (Desikan et al., 2006; Acharya and Assmann, 2009; Zhu et al., 2014). Thus, although the ABA-induced mechanisms regulate both stomata formation and stomatal closure have been widely investigated in *Arabidopsis*, our understanding of how these mechanisms act together with other phytohormones and how they control stomatal movement under stress conditions, where closing and opening stimulus are found simultaneously, remains largely unknown. Given the complexity of these mechanisms, we argue that the only perspective to fully understand

stomatal movement is by using systems biology approaches (Li et al., 2006; Wang et al., 2014b; Medeiros et al., 2015; Medeiros et al., 2018).

Beyond the crosstalk between phytohormones on  $g_s$  regulation, recent evidence indicates a complex regulation of guard cell metabolism and several particularities that differ substantially from mesophyll cells (Santelia and Lawson, 2016). For instance, guard cell  $\text{CO}_2$  metabolism is unique in terms of functioning and regulation. The molecular mechanisms by which guard cells recognize  $\text{CO}_2$  involves a complex network of carbonic anhydrases, kinases such as OST1, HT1, GHR1, and MPK12; as well as anion channels such as SLAC1 and QUAC1 (Engineer et al., 2016a; Hedrich and Geiger, 2017; Negi et al., 2018). Alterations in any of these molecular mechanisms may have significant effects in the stomata response to the environment or under stress. For instance, it was described that mutations in the plasma membrane protein SLAC1 (SLOW ANION CHANNEL ASSOCIATED 1) that mediate guard cell anion efflux by slow anion (S-type) channels altered most of the typical responses to light,  $\text{CO}_2$ , ozone, ABA, humidity,  $\text{Ca}^{2+}$ ,  $\text{H}_2\text{O}_2$  and NO (Vahisalu et al., 2008). Furthermore, higher  $\text{CO}_2$  fixation in the dark and higher PEPc-mediated  $\text{CO}_2$  assimilation have been demonstrated in guard cells of C3 plants, highlighting that the photosynthetic metabolism of these cells does not resemble of those of mesophyll C3 cells (Gotow et al., 1988; Robaina-Estévez et al., 2017). Given the low level of chlorophyll and the few chloroplast numbers found in guard cells (Willmer and Fricker, 1996), RubisCO-mediated  $\text{CO}_2$  assimilation was for long-neglected (Outlaw et al., 1979). However, recent evidence highlights the importance of guard cell chloroplasts for stomatal opening (Wang et al., 2014a; Azoulay-Shemer et al., 2015, 2016). By contrast, guard cells possess a higher mitochondria number and higher respiration rate, when compared to mesophyll cells (Willmer and Friker, 1996; Araújo et al., 2011). These features lead to a unique mode of metabolism which resembles a combination of sink and C4/CAM cells (Cockburn, 1983; Daloso et al., 2016a, 2017).

As a sink cell, guard cells depend on sucrose and other metabolites from source tissues. Evidence supporting this assumption comes also from transcriptomic studies that have shown high expression of sucrose/hexoses transporters found in guard cells and by the higher activity of sink marker enzymes in guard cells (Hite et al., 1993; Ritte et al., 1999; Bates et al., 2012; Bauer et al., 2013a; Daloso et al., 2015). Furthermore, photosynthesis-derived sucrose has been pointed as an important mesophyll to guard cell connectors. Whilst the over accumulation of sucrose and other sugars induce stomatal closure (Kelly et al., 2013; Gago et al., 2016; Li et al., 2016; Medeiros et al., 2018), antisense inhibition of a sucrose importer to guard cells leads to lower guard cell sucrose content and lower  $g_s$  (Antunes et al., 2017b). This indicates that sugars are important metabolites for the A- $g_s$  trade-off regulation (Daloso et al., 2016a; Granot and Kelly, 2019). Indeed, alteration of a subsidiary cell-localized glucose transporter, a SWEET transporter that is regulated by the amount of photoassimilates, substantially alter  $g_s$  (Wang et al., 2019). Thus, it seems likely that sugars, especially sucrose, mediate stomatal closure in periods of high A, as a mechanism to improve WUE (Kang et al., 2007; Daloso et al., 2016a).

Not only sucrose but also mesophyll-derived malate has also been closely associated to the regulation of stomatal movements (Hedrich and Marten, 1993; Talbott and Zeiger, 1993; Laporte et al., 2002; Nunes-Nesi et al., 2007; Araújo et al., 2011; Penfield et al., 2012; Medeiros et al., 2016, 2017). Furthermore, compelling evidence indicates that guard cell malate metabolism does not only affect  $g_s$  but also  $g_m$  (Medeiros et al., 2016), suggesting that both  $g_s$  and  $g_m$  may have common regulation points. Indeed, both  $g_s$  and  $g_m$  have been shown to be couple in a wide range of conditions (Flexas et al., 2002; Flexas et al., 2013a, 2013b). Thus, it is clear that both  $g_s$  and  $g_m$  respond directly to changes in the environment, which affects A. In turn, the  $\text{CO}_2$  assimilated by mesophyll cells can be converted into molecules that will be transported to guard cells where it induces stomatal closure (Lu et al., 1995, 1997; Mott et al., 2008; Mott, 2009). This complex interplay among A,  $g_s$  and  $g_m$  is extremely important in a fluctuating

environment, in which opening and closing environmental signals are combined with (a)biotic interactions in the continuum soil-plant-atmosphere to regulate the  $\text{CO}_2$  flux through the leaf and, consequently, overall plant WUE.

#### 4. Molecular and physiological mechanisms driving mesophyll conductance

Mesophyll conductance to  $\text{CO}_2$  ( $g_m$ ) refers to the facility offered by the leaf for  $\text{CO}_2$  diffusion from the sub-stomatal cavity to the chloroplast stroma where the  $\text{CO}_2$  fixation occurs. This is a complex pathway, comprising aerial, aqueous and lipid phases and composed of several steps each one representing a partial component of  $g_m$ : intercellular air spaces ( $g_{ias}$ ), cell walls ( $g_{cw}$ ), plasma membranes ( $g_{pm}$ ), cytosol ( $g_{cyt}$ ), chloroplast membranes ( $g_{chl}$ ) and stroma ( $g_{st}$ ). Contrary to stomatal conductance, the regulatory pathways of  $g_m$  are poorly understood at the biochemical and molecular levels (Fig. 1).

$g_m$  can be estimated by combining its partial conductance components, which in turn can be approximated dividing values for the diffusivity of  $\text{CO}_2$  at a given temperature in each medium (air, water or lipid) by the distance represented by each step pathway (Evans et al., 2009). For instance,  $g_{cw}$  is calculated considering  $\text{CO}_2$  diffusivity in water and cell wall (CW) thickness (Evans et al., 2009). This is the basis of the widely-used anatomical method to estimate  $g_m$  (Niinemets and Reichstein, 2003; Tosens et al., 2012a; Tomás et al., 2013; Carriqui et al., 2019). However, this is a simplistic view as it assumes that each partial step is composed of a single medium. Hence, this might be unrealistic as, for instance, cell membranes are formed by lipids but also by abundant proteins (Kjellbom and Larsson, 1984; Engelman, 2005); cytosol and stroma are far from being 'pure water' and, although CW pores might be filled mostly of water, chemical interactions between water, apoplast and CW components (including proteins and phenols) (Mercado et al., 2004; Schiraldi et al., 2012) and diffusing  $\text{CO}_2$  should be expected and thus affect the  $\text{CO}_2$  passage through the CW. In addition, the presence of active facilitating mechanisms in any of the step pathways would affect the 'apparent' diffusivity of such pathways. Indeed, while the anatomical method for estimating  $g_m$  often correlates well with *in vivo* estimates of  $g_m$  (e.g. using combined gas exchange and chlorophyll fluorescence or isotope discrimination; see e.g. Pons et al., 2009 and Flexas et al., 2012, 2018a, 2018b for an ample description of the most commonly used methods) in non-stressed plants, the correlation frequently deviates from the 1:1 relationship (Tomás et al., 2013; Fini et al., 2016; Tosens et al., 2016; Peguero-Pina et al., 2017a, b; Veromann-Jürgenson et al., 2017; Carriqui et al., 2019). Moreover,  $g_m$  determined from anatomy often differs strongly from  $g_m$  estimated *in vivo* in plants subject to abiotic stresses, as the former often remains unaffected while abiotic stresses generally induce significant decreases of  $g_m$  (Tomás et al., 2014, but see also Tosens et al., 2012a). This fact – if not an artifact (e.g. Carriqui et al., 2019a) – most likely reflects stress-induced changes in the partial pathways composition and / or in the activity of potential facilitating mechanisms (Flexas and Díaz-Espejo, 2015). However, as said, the regulatory effects of partial pathway compositions and facilitating mechanisms on  $g_m$  are poorly known. Among the former, effects of CW composition on  $g_m$  have been proposed (Niinemets et al., 2009; Gago et al., 2016; Ellsworth et al., 2018; Clemente-Moreno et al., 2019; Roig-Oliver et al., 2020) among potential facilitating mechanisms, chloroplast movements, aquaporins and carbonic anhydrases are the most frequently-invoked candidates (Bernacchi et al., 2002; Terashima et al., 2011; Tomás et al., 2013). Nevertheless, it is likely that the composition of other steps in the  $g_m$  pathway – such as the specific lipid composition of membranes and/or the abundance and typology of membrane-bound or membrane-integral proteins, or the solutes of cytosol and stroma – also have relevant effects on  $\text{CO}_2$  diffusion and hence  $g_m$ , yet none of these have been investigated yet.

#### 4.1. Cell wall dynamics

The structure and interaction of cellulose, hemicelluloses, pectins, lignin, phenols and structural proteins will determine the size of the free spaces within the CW (Carpita et al., 1979; Baron-Epel et al., 1988; Fleischer et al., 1999; Lampert and Kieliszewski, 2005) that the CO<sub>2</sub> should cross in its way to the stroma carboxylation sites (Gago et al., 2019). CWs are dynamic structures that can quickly respond to their environment changing both chemical composition and physical structure (Houston et al., 2016). For instance, under salt or drought stress, the apoplastic media that is usually acidic, increase their pH which may difficulting the CO<sub>2</sub> solubilization and alter proteins functionality. Moreover, in these stress conditions, the oxidative response includes both ROS increase and the activation of different apoplastic enzymes that are also key players in CW remodelling (Pitann et al., 2009; O'Brien et al., 2012; Tenhaken, 2015). Moreover, it is known that apoplastic ROS levels are major signalling components (as it is also known for guard cell movements as described above) for proteins in charge of the communication between the extracellular environment to the intracellular compartment (Pignocchi and Foyer, 2003), that should drive subsequently highly regulated molecular mechanisms and metabolic fluxes responses through the CW (Verbančič et al., 2018).

Antioxidant apoplastic enzymes as peroxidases (POXs) or superoxide dismutases (SODs) can act as stiffening agents catalysing the cross-linking between different CW matrix compounds including phenolic groups and lignification (Wu and Cosgrove, 2000). For instance, even short-term drought frequently increases lignification of CWs (Escudero et al., 1992; Sobrado, 1992; Henry et al., 2000) and has been suggested to decrease  $g_{cw}$  due to the occlusion of pores (Monties, 1989; Niinemets et al., 2009). As well, POXs can also drive the CW loosening via oxidative cleavage of other polymers that show important hydrocolloid properties to retain water as xyloglucan and pectins, thus affecting CW composition and porosity (Cathala and Monties, 2001; Panchev et al., 2010; Tenhaken, 2015; Schmidt et al., 2016). Interestingly, ascorbate, the most abundant non-enzymatic antioxidant in the apoplast, and the symplastic ascorbate oxidase activity might regulate POXs activity controlling hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) levels (Ros-Barceló et al., 2006) and could have a regulatory effect on POX-CW remodelling, indicating the complex network regulation between apoplast and symplast. Additional line of evidence can be observed in the CW profile of angiosperm resurrection species that show higher levels of hemicelluloses, pectines and structural proteins (pectin-arabinans, arabinogalactan proteins and arabinoxylans) than other non-tolerant desiccation angiosperm species. This exclusive profile is related to highly flexible CW that can support significant mechanical pressures without collapse, plus subsequently, these compounds facilitate rehydration after a desiccation event (Moore et al., 2013).

Interestingly, significant associations of CW compounds were described including, for example, galactose, xylose and arabinose (precursors of pectins and hemicellulose) and the shikimate pathway through hydroxybenzoate (precursor of several phenolic compounds) relate specifically with  $g_m$  but were mostly not observed for  $g_s$  or  $A_n$  through a multi-species modelling analysis. Additional information was obtained because  $g_m$  associations showed a higher number of amino acids than  $g_s$  and  $A$  (Gago et al., 2016), being speculated if these associations can refer to the activity and levels of aquaporins, carbonic anhydrases and structural CW proteins. These data indirectly signalled that the intrinsic CW composition could directly affect the CO<sub>2</sub> pathway beyond the known inverse anatomical relationship with its thickness (Gago et al., 2019). Lately, Ellsworth et al. (2018) also observed a reduction of  $g_m$  in mutants with disrupted synthesis and accumulation of the hemicelluloses (1,3)- and (1,4)-linked  $\beta$ -glucosyl polysaccharides. Further evidence was found recently, showing that  $g_m$  was related to the peroxidase activity at the apoplast level but not at the symplast level in response to salt and drought stress (and its recovery) (Clemente-Moreno et al., 2019). In parallel,  $g_m$  reductions were also related significantly

with pectins increase and positively with the ratio hemicelluloses/pectins, and again with specific CW primary metabolites as galactose (pectins precursor), GABA, glucosamine, and *p*-coumarate (phenolic compound); altogether, indicating a complex response affecting  $g_m$  by oxidative stress and redox status driving CW remodeling.

Further research is needed to disentangle if the polymer hydrocolloids accumulation (hemicelluloses, pectins and phenols) can enlarge CW lattice (thus, increase the CO<sub>2</sub> pathway) or these increases in the hydrophilic lattice would facilitate CO<sub>2</sub> solubilization and movement across CW under stress. In this sense, the unknown role of the structural proteins remains as much challenging as exciting. These studies provided the first evidence that CW composition affects  $g_m$ , yet the complete mechanisms remain mostly unknown.

#### 4.2. Chloroplast movements

Chloroplasts are generally as close to CWs as possible in order to short the liquid pathway of CO<sub>2</sub> through cytosol – maximize  $g_{cyt}$  (Evans et al., 2009; Flexas et al., 2018a, 2018b). In fact, the chloroplast surface area facing intercellular air spaces per unit leaf area ( $S_c/S$ ) is one of the anatomical parameters that explain more variance of  $g_m$  in studies of non-stressed diverse species (Syvertsen et al., 1995; Hanba et al., 2002; Carriqui et al., 2015; Tosens et al., 2016; Veromann-Jürgenson et al., 2017; Peguero-Pina et al., 2017a, b; Carriqui et al., 2019). This parameter also changes under drought, sunlight and potassium treatments (Tosens et al., 2012a, 2012b; Fini et al., 2016; Lu et al., 2016; Han et al., 2019), with changes in leaf development (Miyazawa and Terashima, 2001), as well as in some mutants and transgenic plants with altered chloroplast shapes, numbers or distribution (Sharkey et al., 1991; Evans et al., 1994; Weraduwage et al., 2018). Thus, some authors have suggested that changes in  $S_c/S$  could explain the short-term responses of  $g_m$  e.g. to light (Ho et al., 2016; Thérout-Rancourt and Gilbert, 2017) and high temperatures (Flexas and Díaz-Espejo, 2015). However, several studies have observed scarce changes of  $g_m$  associated with chloroplast movement (Gorton et al., 2003; Loreto et al., 2009; Carriqui et al., 2019a) and only Tholen et al. (2008) observed a correlation between short-term variations of  $g_m$  in response to light quality with changes in chloroplast placement and  $S_c/S$ . The involvement of chloroplasts' movements on  $g_m$  was further corroborated using phytochrome-mutants lacking chloroplast movements (Tholen et al., 2008). Therefore, more pieces of evidence are needed to associate the short-term decrease of  $g_m$  with fast removal of chloroplast from the outer cell surface.

#### 4.3. Aquaporins

During the last decades, the importance of aquaporins as facilitators of the diffusion of CO<sub>2</sub> across membranes has been identified (reviewed in Uehlein et al., 2017 and Flexas et al., 2016). First, Terashima and Ono (2002) demonstrated that  $g_m$  decreased after the application of HgCl<sub>2</sub>, a general inhibitor of aquaporins. Subsequently, the implication of tobacco NtAQP1 aquaporin in the permeabilization for CO<sub>2</sub> of plasma membranes (Uehlein et al., 2003; Otto et al., 2010) and chloroplast membranes (Uehlein et al., 2008) was evidenced. Genetically-modified plants had been also used to verify *in planta* the involvement of NtAQP1 (Flexas et al., 2006; Uehlein et al., 2008; Sade et al., 2014) and some other aquaporins (Hanba et al., 2004; Heckwolf et al., 2011; Kawase et al., 2013) on  $g_m$ . While all those studies were performed in non-stressed plants, Miyazawa et al. (2008) suggested by treating irrigated and water-stressed plants with HgCl<sub>2</sub> a specific involvement of aquaporins in drought-induced decreases of  $g_m$ . A similar suggestion was made by Perez-Martin et al. (2014) after monitoring aquaporin transcription in water-stressed plants. Despite of all these proofs, recently Kromdijk et al., 2019 have shown a null effect of aquaporins knock-out on  $g_m$  measured with multiple methods. One of the explanations suggested by these authors is that redundancy of different aquaporins isoforms would mask the effects of a single gene knock-out.



**Table 1**

Coordination of the stomatal ( $g_s$ ) and mesophyll ( $g_m$ ) conductances short-term responses (min-hours) to several environmental factors and ABA application. The responses are highlighted in blue when  $g_s$  and  $g_m$  display the same trend and in red when they differ. <sup>a</sup> Only high nitrogen plants included. <sup>b</sup> Lines studied were CML333, IL14H, Ki11, Mo17 and W22; all five presented the same  $g_s$  and  $g_m$  response.

| Treatment                                                               | Species                          | Genotype/line        | $g_s$     | $g_m$ | Reference                  |
|-------------------------------------------------------------------------|----------------------------------|----------------------|-----------|-------|----------------------------|
| Low CO <sub>2</sub><br>(C <sub>a</sub> < 400 ppm)                       | <i>Triticum aestivum</i>         | Yecora 70            | ↓         | =     | Tazoe et al. (2009)        |
|                                                                         | <i>Eucalyptus globulus</i>       |                      | ↑         | ↑     | Douthe et al. (2011)       |
|                                                                         | <i>Eucalyptus saligna</i>        |                      | ↑         | ↑     | Douthe et al. (2011)       |
|                                                                         | <i>Eucalyptus sieberi</i>        |                      | ↑         | ↑     | Douthe et al. (2011)       |
|                                                                         | <i>Oryza sativa</i>              | Heshengwanyou        | ↑         | ↑     | Xiong et al., 2015,        |
| High CO <sub>2</sub><br>(C <sub>a</sub> > 400 ppm)                      | <i>Zea mays</i>                  | Various <sup>b</sup> | ↑         | ↑     | Kolbe et al., 2018         |
|                                                                         | <i>Nicotiana tabacum</i>         | Samsun               | ↓         | ↓     | Flexas et al. (2007)       |
|                                                                         | <i>Nicotiana sylvestris</i>      |                      | ↓         | ↓     | Flexas et al. (2007)       |
|                                                                         | <i>Triticum aestivum</i>         | Yecora 70            | =         | =     | Tazoe et al. (2009)        |
|                                                                         | <i>Eucalyptus globulus</i>       |                      | =         | =     | Douthe et al. (2011)       |
|                                                                         | <i>Eucalyptus saligna</i>        |                      | =         | =     | Douthe et al. (2011)       |
|                                                                         | <i>Eucalyptus sieberi</i>        |                      | ↓         | =     | Douthe et al. (2011)       |
|                                                                         | <i>Oryza sativa</i>              | Heshengwanyou        | ↓         | ↓     | Xiong et al., 2015,        |
|                                                                         | <i>Zea mays</i>                  | Various <sup>b</sup> | ↓         | =     | Kolbe et al., 2018         |
|                                                                         | <i>Arabidopsis thaliana</i>      | Col-0                | ↓         | ↓     | Mizokami et al. (2019)     |
| Low light<br>(PPFD < 4[MNN13] 00 μmol m <sup>-2</sup> s <sup>-1</sup> ) | <i>Arabidopsis thaliana</i>      | ost1                 | =         | ↓     | Mizokami et al. (2019)     |
|                                                                         | <i>Arabidopsis thaliana</i>      | sac1 – 2             | =         | ↓     | Mizokami et al. (2019)     |
|                                                                         | <i>Triticum aestivum</i>         | Yecora 70            | = [MNN12] | =     | Tazoe et al. (2009)        |
|                                                                         | <i>Eucalyptus globulus</i>       |                      | ↓ [MNN14] | ↓     | Douthe et al. (2011, 2012) |
|                                                                         | <i>Eucalyptus saligna</i>        |                      | ↓         | ↓     | Douthe et al. (2011, 2012) |
| High light<br>(PPFD > 1000 μmol m <sup>-2</sup> s <sup>-1</sup> )       | <i>Eucalyptus sieberi</i>        |                      | ↓         | ↓     | Douthe et al. (2011, 2012) |
|                                                                         | <i>Triticum aestivum</i>         | Yecora 70            | ↑ [MNN16] | =     | Tazoe et al. (2009)        |
|                                                                         | <i>Eucalyptus globulus</i>       |                      | = [MNN17] | =     | Douthe et al. (2011, 2012) |
|                                                                         | <i>Eucalyptus saligna</i>        |                      | =         | =     | Douthe et al. (2011, 2012) |
|                                                                         | <i>Eucalyptus sieberi</i>        |                      | =         | =     | Douthe et al. (2011, 2012) |
|                                                                         | <i>Oryza sativa</i>              | Heshengwanyou        | ↑ [MNN18] | ↑     | Xiong et al., 2015,        |
|                                                                         | <i>Eucalyptus tereticornis</i>   |                      | ↑ [MNN19] | ↑     | Campany et al. (2016)      |
|                                                                         | <i>Phlebotidium aureum</i>       |                      | = [MNN20] | =     | Xiong et al. (2018)        |
|                                                                         | <i>Nephrolepis cordifolia</i>    |                      | =         | =     | Xiong et al. (2018)        |
|                                                                         | <i>Taxus baccata</i>             |                      | ↑         | ↑     | Xiong et al. (2018)        |
| B[MNN21] lue light                                                      | <i>Ginkgo biloba</i>             |                      | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Nerium oleander</i>           |                      | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Populus nigra</i>             |                      | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Gossypium hirsutum</i>        |                      | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Helianthus annuus</i>         |                      | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Centella asiatica</i>         |                      | =         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Oryza sativa</i>              | Shanyou 63           | ↑         | ↑     | Xiong et al. (2018)        |
|                                                                         | <i>Arabidopsis thaliana</i>      | Col                  | ↑         | ↓     | Tholen et al. (2008)       |
|                                                                         | <i>Arabidopsis thaliana</i>      | gl1                  | ↑         | ↓     | Tholen et al. (2008)       |
|                                                                         | <i>Arabidopsis thaliana</i>      | phot2                | ↑         | =     | Tholen et al. (2008)       |
| ABA application                                                         | <i>Arabidopsis thaliana</i>      | chup1                | ↑         | =     | Tholen et al. (2008)       |
|                                                                         | <i>Nicotiana plumbaginifolia</i> | WT                   | ↓         | ↓     | Mizokami et al., 2015      |
|                                                                         | <i>Nicotiana plumbaginifolia</i> | aba1                 | ↓         | ↓     | Mizokami et al., 2015      |
|                                                                         | <i>Prunus avium</i>              |                      | ↓         | ↓     | Sorrentino et al., 2016    |
|                                                                         | <i>Populus nigra</i>             |                      | ↓         | ↓     | Sorrentino et al., 2016    |
|                                                                         | <i>Olea europaea</i>             |                      | ↓         | ↓     | Sorrentino et al., 2016    |
|                                                                         | <i>Rosoidea rosa</i>             |                      | ↓         | ↓     | Sorrentino et al., 2016    |
|                                                                         | <i>Arabidopsis thaliana</i>      | Camp David           | ↓         | ↓     | Sorrentino et al., 2016    |
|                                                                         | <i>Arabidopsis thaliana</i>      | Col-0                | ↓         | ↓     | Mizokami et al. (2019)     |
|                                                                         | <i>Arabidopsis thaliana</i>      | ost1                 | =         | =     | Mizokami et al. (2019)     |
|                                                                         | <i>Arabidopsis thaliana</i>      | sac1 – 2             | ↓         | =     | Mizokami et al. (2019)     |

The number of aquaporins belonging to 10 different aquaporin families and multiple isoforms present in each species is generally unknown (Kjellbom et al., 1999; Maurel et al., 2008) and the CO<sub>2</sub> transport activity may differ among aquaporin types (Otto et al., 2010). Furthermore, the role that aquaporins play in the short-term response of  $g_m$  to temperature, CO<sub>2</sub> and light (Bernacchi et al., 2002; Flexas et al., 2007; Xiong et al., 2015,) is not fully understood yet.

#### 4.4. Carbonic anhydrases

Carbonic anhydrases (CAs) are cytosol, mitochondrial and stromal enzymes that catalyse the reversible reaction  $\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{HCO}_3^- + \text{H}^+$  to allow a suitable diffuse through the liquid phase and a sustained CO<sub>2</sub> supply at the carboxylation site of Rubisco. It is under debate if CAs have an influence on the changes of  $g_m$  or whether CO<sub>2</sub> interconversion is not a limiting factor of  $g_m$  (Tholen and Zhu, 2011; Flexas et al., 2012; Veromann-Jürgenson et al., 2017). In C<sub>4</sub> plants,

photosynthetic activity and  $g_m$  seemed reliant on the activity of CAs by assisting the high CO<sub>2</sub> concentration of chloroplast of sheath cells (Badger and Pfanz, 1995; Ludwig, 2012; Studer et al., 2014; Osborn et al., 2017; Kolbe et al., 2018; but see also Cousins et al., 2006). In C<sub>3</sub> plants, pieces of evidence are not as clear. Variation of CAs expression or activity by the use of inhibitors, mutants or water stress has produced inconclusive results, as studies have reported either an effect on  $g_m$  (Badger and Pfanz, 1995; Gillon and Yakir, 2000; Perez-Martin et al., 2014; Momayyezi and Guy, 2017a, b; Momayyezi and Guy, 2018) or a null effect (Price et al., 1994; Williams et al., 1996; Sasaki et al., 1998). Thus, the role of CAs in determining  $g_m$  remains a matter of debate (Momayyezi et al., 2020), as well as, the relative importance of the biochemical limitations to anatomical determinants of  $g_m$ .



## 5. The molecular and physiological inter-play between both conductances and photosynthesis

It is widely described that both  $g_s$  and  $g_m$  show a notable coupled behaviour largely influencing photosynthesis and WUE (Flexas et al., 2013a, 2013b; Gago et al., 2014). There is a significant variation in the  $g_m / g_s$  ratio across species and even genotypes (Flexas et al., 2013a, 2013b), which can be an important feature for WUE selection, especially under drought conditions (Galmés et al., 2011; Thérroux-Rancourt et al., 2015). An increased  $g_m / g_s$  ratio (i.e. a decreased mesophyll conductance limitation to photosynthesis) seems to be also at the basis of the observed progressive increase in WUE along land plant's phylogeny, from lowest values in bryophytes to the largest and more variable in angiosperms (Flexas and Carriqui, 2020). In fact, stomatal density and regulation have been often genetically manipulated and resulted in plants with improved WUE (Antunes et al., 2012; Kelly et al., 2013; Wang et al., 2014c; Medeiros et al., 2016; Daloso et al., 2016b; Antunes et al., 2017b; Hughes et al., 2017; Vráblová et al., 2017). Recently, it was demonstrated in tobacco plants the effect of one macronutrient, the chloride ( $Cl^-$ ), prompted higher WUE driven by lower  $g_s$  and higher  $g_m$ , increasing  $g_m / g_s$  ratio without penalizing A. These trends were driven by anatomical traits at stomata level as reducing stomata density but increasing their size, in combination with an elevated  $S_c / S$  (higher number of smaller chloroplast facing the mesophyll airspaces) leading to higher  $g_m$  (Franco-Navarro et al., 2019).

Hence, both conductances, although coordinated when considering large interspecific datasets, show a certain degree of uncoupling under short-term conditions, which may arise from their different underlying mechanisms (Gallé et al., 2011; Flexas et al., 2012; Clemente-Moreno et al., 2019). The response of  $g_m$  and  $g_s$  to water and salt stresses has already been extensively reviewed elsewhere (Flexas et al., 2013a, 2018a, 2018b; Nadal and Flexas, 2019), but the response of  $g_m$  and its potential drivers to well-known stimuli affecting stomatal response are still poorly understood.

We compiled a multi-species dataset from 13 papers published in the literature covering in 23 species the responses of  $g_s$  and  $g_m$  to short-term changes in atmospheric  $CO_2$ , light irradiances as well exogenous ABA application (Table 1). Notably, in most cases  $g_s$  and  $g_m$  show a similar trend, indicating a rather conserved response to stimuli and directly affecting photosynthesis (light intensity,  $CO_2$ ). This positive coordination may arise from the need to reduce water losses and maximize  $CO_2$  availability for photosynthesis during limited optimum conditions, as for example in the case of sunflecks in understorey vegetation (Campany et al., 2016).

The short-term response of  $g_m$  to both high light and  $CO_2$  was first reported by Flexas et al. (2007), where  $g_m$  showed a rapid increase under low  $CO_2$  (< 400 ppm) and a decrease under high  $CO_2$  (> 400 ppm) (Table 1), in a similar manner as the reported response of  $g_s$  in angiosperms (Brodrick et al., 2009). However, this pattern may partially arise from mathematical circularity due to the inclusion of  $C_i$  in the methods used for  $g_m$  estimation (Gu and Sun, 2014a, 2014b). The lack of an anatomical basis for the apparent variation of  $g_m$  with  $CO_2$  adds to the potential artifactual nature of this response (Carriqui et al., 2019a) and the lack of good understanding of the molecular factors that could drive quick responses of  $g_m$  (Shatil-Cohen et al., 2011; Grondin et al., 2015; Gago et al., 2016; Clemente-Moreno et al., 2019; Mizokami et al., 2019) avoids further concluding on this aspect of  $g_m$  regulation, a topic that clearly deserves further investigation. Similarly,  $g_m$  has been reported to vary with light intensity (Douthe et al., 2011, 2012) (Table 1), which again may partially arise from model limitations (Gu and Sun, 2014a, 2014b). Nonetheless, the fact that the response of  $g_m$  to light and  $CO_2$  is not universal across all species (Xiong et al., 2018) seems to indicate that part of the observed variation is not entirely artifactual but responds to some species-specific features. In this case, leaf anatomy may play an important role as the apparent response of  $g_m$

to light intensity would emerge from the differential light penetration in the mesophyll layers, instead of any biochemistry-driven process (Thérroux-Rancourt and Gilbert, 2017). On the other hand, the response of  $g_m$  to blue light is well supported by changes in  $S_c$  due to chloroplast re-arrangements (Tholen et al., 2008). Interestingly, blue light is one of the few cases where  $g_m$  and  $g_s$  show an opposite response (Table 1).

During drought, the decline of  $g_m$  and  $g_s$  could be partially attributed to a hydraulic ( $K_{leaf}$ ) decline (Rodríguez-Domínguez et al., 2016; Wang et al., 2018), although the interplay between these parameters is still under debate (Flexas et al., 2018a, 2018b). Interestingly, at the leaves, ABA is mostly produced in the mesophyll cells (McAdam and Brodrick, 2018) and its action in the guard cells are highly regulated by several compounds from ROS levels, hormones, antioxidants and peptides produced in another tissues (Watkins et al., 2014; Takahashi et al., 2018). In general, ABA is considered an important factor in the decline of both conductances. For example, exogenous ABA application promptly decreases to a similar extent  $g_m$  and  $g_s$  in several species (Vrábl et al., 2009; Mizokami et al., 2015, 2019; Sorrentino et al., 2016) (Table 1). However, in poplar, the sensitivity of  $g_m$  to ABA is probably lower than that of  $g_s$ , as  $g_s$  in well-watered poplar changed concomitantly with accordingly with slight daily variations in ABA but  $g_m$  did not (Brunetti et al., 2019). This may indicate a different pathway for ABA sensitivity, which is also suggested in *Arabidopsis* mutant lines lacking OST1 and SAC1, where the  $g_m$ - $g_s$  coupling is hindered (Mizokami et al., 2019). Aquaporins could be responsible for the ABA-mediated  $g_m$  decrease. ABA mediates decreases in membrane water permeability via aquaporin activity down-regulation, resulting in a reduction in  $K_{leaf}$  (Shatil-Cohen et al., 2011). Indeed, exogenous ABA application affects the expression of several aquaporins members of the PIP subgroup in *Arabidopsis* (Jang et al., 2004). Recently, light has been brought in this matter, it was published that stomata closure required the activity of the PIP2;1, especially in response to ABA (increases osmotic water permeability in guard cells) in *Arabidopsis* (Grondin et al., 2015). Interestingly, authors observed that normal stomata functionality was recovered by the overexpression of Open stomata 1 (OST1) / Snf1-related protein kinase 2.6 (SnRK2.6) (a kinase involved in guard cell ABA signalling) in the knock-out plants mediated by a phosphorylation mechanism in the cytosolic PIP2;1 peptide at Ser-121.

Further connections can be established also with the role of CAs related to the interplay between both conductances. While the role of CAs in  $g_m$  has been discussed in previous sections and recently reviewed (Momayyezi et al., 2020), additionally *Arabidopsis* lacking several CAs showed limited  $g_s$  response to  $CO_2$  changing concentrations, higher  $g_s$  and density compared to wild type (Hu et al., 2010, 2015; Engineer et al., 2014). On the other hand, it has been also suggested that CAs can act as  $CO_2$  sensors for guard cells supplying  $HCO_3^-$  for OST1 kinase that regulates S-type anion channels related to  $CO_2$  stomata closure (Hu et al., 2010, 2015; Xue et al., 2011).

In the previous section, important differences in stomatal responsiveness to abiotic factors among phylogenetic plant groups has been highlighted. In this sense given that mesophyll photosynthetic capacity differs substantially between angiosperms and basal lineage of plants (Tosens et al., 2016; Gago et al., 2019), it is reasonable to hypothesize that mesophyll metabolism may play a central role in the regulation of stomatal movements across plant evolution. This idea is strengthened by the fact that guard cell metabolism cannot sustain stomatal movements by itself (Messinger, 2006; Mott, 2009), but depending, from mesophyll-derived metabolites such as sugars and organic acids that are charged into the apoplast and transported to guard cells through the transpiration stream (Fujita et al., 2013; Nunes-Nesi et al., 2011; Daloso et al., 2016; Santelia and Lawson, 2016; Daloso et al., 2018). What still remains unknown is whether and to which extent signals from mesophyll cells contribute to the differential velocity of stomatal responses observed among angiosperms and other plant groups, and how it is possible to use this knowledge to optimize water use efficiency.

It seems reasonable to speculate that another possible link between

both conductances is the redox status of leaves and the whole plant. ROS were traditionally considered as toxic byproducts of aerobic metabolism, i.e. photosynthesis, and cells should deal with them to avoid excessive oxidation and irreversible damage; however, currently, ROS are being recognized as essential signalling molecules for several processes regulation (Mittler, 2017; Foyer, 2018). In agreement, evidence suggesting the association of NAD<sup>+</sup> metabolism with stomatal function (Hashida et al., 2010), as well as the NAD<sup>+</sup> transport between organelles regulating guard cell biogenesis have been found in *Arabidopsis* (de Souza Chaves et al., 2019d). For instance, it was discovered the contribution of the aquaporin AtPIP2;1 to the intracellular accumulation of H<sub>2</sub>O<sub>2</sub> (facilitating water efflux required for stomatal closure), that is also related with superoxide dismutases and CW peroxidases that drive the production of apoplastic ROS and CW remodelling (Wu and Cosgrove, 2000; Khokon et al., 2011; Kadota et al., 2014; Rodrigues et al., 2017). ROS can not only alter  $g_s$ , but also major described components of  $g_m$  as the functionality of diverse enzymes related to  $g_m$  as AQPs and CAs (see sections 4.3 and 4.4), cell and chloroplast membranes lipidic profile (changes in the lipid species, saturation and length of their chains) that will alter their permeability and the activity of transmembrane proteins as AQPs (Lee, 2004; Phillips et al., 2009a, 2009b; Burgos et al., 2011) and drive major changes into the CW composition (Tenhaken, 2015; Clemente-Moreno et al., 2019). Under stress conditions, ROS signaling and process regulation can become even more essential (before oxidative burst will promote physiological damage), and thus the altered redox status can drive a pool of mechanisms in order to deal with abiotic stress.

## 6. Concluding remarks

Now, global food security is a major worldwide concern attempting to the predicted climate change scenarios for the most important agricultural regions on the planet and the growing human population. The current food and fuel requirements are obtained based on large water consumption and nutrients, is the new challenge for society and the scientific community to ensure or even increase productivity in these predicted scenarios.

Diffusional limitations (i.e.  $g_s$  and  $g_m$ ) to photosynthesis can impose easily about two-thirds of the total photosynthetic limitation under optimum conditions, but under mild and severe abiotic stress conditions they can reach > 85 % for most of the studied species already characterized (Flexas et al., 2018a, 2018b; Gago et al., 2019). Thus, knowledge about both conductances is essential to improve WUE, targeted crop breeding programs and sustainable, resource-efficient and environmental-friendly agriculture strategies. Curiously, the knowledge acquired from decades of research from  $g_s$  and  $g_m$  mainly differs in their origin. Biochemical regulation and signaling pathways are highly defined for  $g_s$  and there is still a lot of controversy regarding the molecular mechanisms, however, an important lack of knowledge is observed for its structural determinants. The opposite is true for  $g_m$ , known as a highly-complex parameter, anatomical determinants are much more well-characterized than its biochemical regulation and signaling pathways that still remains mostly unknown. Even more scarce is the information in the literature discussing the known coupled behaviour that frequently shows both conductances under most of the conditions (Flexas et al., 2013a) and, moreover, establishing the mechanisms that could be the common determinants linking their dynamics.

Given the complexity of this challenging research focus, integrative multi-disciplinary studies are needed to join ecophysiology with molecular biology using systems biology approaches (Gago et al., 2016; Flexas and Gago, 2018; Medeiros et al., 2019). What remains as key targets for the future would be 1) to complete the mentioned lack of information for each of the conductances, 2) to determine the putative common molecular mechanisms driving the coupled behaviour between both conductances and 3) to use this knowledge for improving

crop breeding strategies to increase maximal photosynthetic capacity and WUE in the global change scenario.

## Author Statement

J.G., D.M.D and J.F. conceived and designed the idea of this review and wrote the first draft of the paper with subsequent inputs from all co-authors. M.N. developed the dataset compilation. D.M.D, W.L.A, A.N.N. drive the writing of Section 3. J.F, M.N., A.V.P.C., M.C.M, M.C develop Section 4. M.M, M.C.M. and J.G. lead Section 5. M.M., J.G., and J.F. design the figures.

## Declaration of Competing Interest

The authors declare no conflicts of interest.

## Acknowledgments

J.F. and J.G. acknowledge the funding obtained from the Spanish Ministry of Science and Technology, Project EREMITA, [PGC2018-093824-B-C41]. A.N.N., W.L.A. and D.M.D. also thank the fund support from National Council for Scientific and Technological Development (CNPq-Brazil, Grants 306818/2016-7, 402511/2016-6 and 428192/2018-1), and the FAPEMIG (Foundation for Research Assistance of the Minas Gerais State, Brazil, Grant RED-00053-16), and also their research fellowships financed by the CNPq-Brazil. M.N. is grateful by his predoctoral fellowship BES-2015-072578 from the Spanish Ministry of Science and co-financed by the European Social Fund. M.M. recognized her postdoctoral fellowship FJCI-2016-31007 ('Juan de la Cierva-Formación' program) co-funded by the Spanish Ministry of Science, Innovation and Universities, the State Research Agency and the University of the Balearic Islands.

## References

- Acharya, B.R., Assmann, S.M., 2009. Hormone interactions in stomatal function. *Plant Mol. Biol.* 69, 451–462.
- Ando, E., Kinoshita, T., 2018. Red light-induced phosphorylation of plasma membrane H<sup>+</sup> ATPase in stomatal guard cells. *Plant Physiol.* 178, 838–849.
- Antunes, W.C., Provart, N.J., Williams, T.C.R., Loureiro, M.E., 2012. Changes in stomatal function and water use efficiency in potato plants with altered sucrolytic activity. *Plant Cell Environ.* 35, 747–759.
- Antunes, W.C., Daloso, D.M., Pinheiro, D.P., Williams, T.C.R., Loureiro, M.E., 2017a. Guard cell-specific down-regulation of the sucrose transporter SUT1 leads to improved water use efficiency and reveals the interplay between carbohydrate metabolism and K<sup>+</sup> accumulation in the regulation of stomatal opening. *Environ. Exp. Bot.* 135, 73–85.
- Antunes, W.C., de Menezes Daloso, D., Pinheiro, D.P., Williams, T.C.R., Loureiro, M.E., 2017b. Guard cell-specific down-regulation of the sucrose transporter SUT1 leads to improved water use efficiency and reveals the interplay between carbohydrate metabolism and K<sup>+</sup> accumulation in the regulation of stomatal opening. *Environ. Exp. Bot.* 135, 73–85.
- Araújo, W.L., Nunes-Nesi, A., Osorio, S., Usadel, B., Fuentes, D., Nagy, R., et al., 2011. Antisense inhibition of the iron-sulphur subunit of succinate dehydrogenase enhances photosynthesis and growth in tomato via an organic acid-mediated effect on stomatal aperture. *Plant Cell* 23, 600–627.
- Assmann, S.M., Jegla, T., 2016. Guard cell sensory systems: recent insights on stomatal responses to light, abscisic acid, and CO<sub>2</sub>. *Curr. Opin. Plant Biol.* 33, 157–167.
- Azoulay-Shemer, T., Palomares, A., Bagheri, A., Israelsson-Nordstrom, M., Engineer, C.B., Bargmann, B.O.R., et al., 2015. Guard cell photosynthesis is critical for stomatal turgor production, yet does not directly mediate CO<sub>2</sub> and ABA-induced stomatal closing. *Plant J.* 83, 567–581.
- Azoulay-Shemer, T., Bagheri, A., Wang, C., Palomares, A., Stephan, A.B., Kunz, H.H., Schroeder, J.I., 2016. Starch biosynthesis in guard cells but not in mesophyll cells is involved in CO<sub>2</sub>-induced stomatal closing. *Plant Physiol.* 171, 788–798.
- Badger, M.R., Pfan, H., 1995. Effect of carbonic anhydrase inhibition on photosynthesis by leaf pieces of C3 and C4 plants. *Funct. Plant Biol.* 22, 45–49.
- Baroli, I., Price, G.D., Badger, M.R., von Caemmerer, S., 2007. The contribution of photosynthesis to the red light response of stomatal conductance. *Plant Physiol.* 146, 737–747.
- Baron-Epel, O., Gharyal, P.K., Schindler, M., 1988. Pectins as mediators of wall porosity in soybean cells. *Planta* 175, 389–395.
- Bates, G.W., Rosenthal, D.M., Sun, J., Chattopadhyay, M., Pepper, E., Yang, J., et al., 2012. A comparative study of the *Arabidopsis thaliana* guard-cell transcriptome and its modulation by sucrose. *PLoS One* 7, e49641.
- Bauer, H., Ache, P., Lautner, S., Fromm, J., Hartung, W., Al-Rasheid, K.A.S., et al., 2013a. The stomatal response to reduced relative humidity requires guard cell-autonomous

- ABA synthesis. *Curr. Biol.* 23, 53–57.
- Bauer, H., Ache, P., Wohlfart, F., Al-Rasheid, K.A.S., Sonnewald, S., Sonnewald, U., et al., 2013b. How do stomata sense reductions in atmospheric relative humidity? *Mol. Plant* 6, 1703–1706.
- Bergmann, D., 2006. Stomatal development: from neighborly to global communication. *Curr. Opin. Plant Biol.* 9, 478–483.
- Bernacchi, C.J., Portis, A.R., Nakano, H., von Caemmerer, S., Long, S.P., 2002. Temperature response of mesophyll conductance. Implications for the determination of Rubisco enzyme kinetics and for limitations to photosynthesis *in vivo*. *Plant Physiol.* 130, 1992–1998.
- Bertolino, L.T., Caine, R.S., Gray, J.E., 2019. Impact of stomatal density and morphology on water-use efficiency in a changing world. *Fr. Plant Sci.* 10, 225.
- Björkman, O., Holmgren, P., 1963. Adaptability of the photosynthetic apparatus to light intensity in ecotypes from exposed and shaded habitats. *Physiol. Plant.* 16, 889–914.
- Boyce, C.K., Brodribb, T.J., Feild, T.S., Zwieniecki, M.A., 2009. Angiosperm leaf vein evolution was physiologically and environmentally transformative. *Proc. R. Soc. B.* 276, 1771–1776.
- Brodribb, T.J., Buckley, T.N., 2018. Leaf water transport: a core system in the evolution and physiology of photosynthesis. *The Leaf: A Platform for Performing Photosynthesis*. Springer, Cham, pp. 81–96.
- Brodribb, T.J., Feild, T.S., 2000. Stem hydraulic supply is linked to leaf photosynthetic capacity: evidence from New Caledonian and Tasmanian rainforests. *Plant Cell Environ.* 23, 1381–1388.
- Brodribb, T.J., Holbrook, N.M., 2004. Stomatal protection against hydraulic failure: a comparison of coexisting ferns and angiosperms. *New Phytol.* 162, 663–670.
- Brodribb, T.J., McAdam, S.A.M., 2011. Passive origins of stomatal control in vascular plants. *Science* 331, 582–585.
- Brodribb, T.J., McAdam, S.A.M., 2013. Unique responsiveness of angiosperm stomata to elevated CO<sub>2</sub> explained by calcium signalling. *PLoS One* 8, 1–13.
- Brodribb, T.J., McAdam, S.A., 2017. Evolution of the stomatal regulation of plant water content. *Plant Physiol.* 174, 639–649.
- Brodribb, T.J., Feild, T.S., Jordan, G.J., 2007. Leaf maximum photosynthetic rate and venation are linked by hydraulics. *Plant Phys.* 144 (4), 1890–1898.
- Brodribb, T.J., McAdam, S.A.M., Jordan, G.J., Feild, T.S., 2009. Evolution of stomatal responsiveness to CO<sub>2</sub> and optimization of water-use efficiency among land plants. *New Phytol.* 183 (3), 839–847.
- Brodribb, T.J., Sussmilch, F., McAdam, S.A.M., 2020. From reproduction to production, stomata are the master regulators. *Plant J.* 101, 756–767.
- Brunetti, C., Gori, A., Marino, G., Latini, P., Sobolev, A.P., Nardini, A., Haworth, M., Giovannelli, A., Capitani, D., Loreto, F., Taylor, G., Mugnozza, G.S., Harfouche, A., Centritto, M., 2019. Dynamic changes in ABA content in water-stressed *Populus nigra*: effects on carbon fixation and soluble carbohydrates. *Ann. Bot.* 124 (4), 627–643.
- Buckley, T.N., 2019. How do stomata respond to water status? *New Phytol.* 224, 21–36.
- Burgos, A., Szymanski, J., Seiwert, B., Degenkolbe, T., Hannah, M.A., Giavalisco, P., Willmitzer, L., 2011. Analysis of short-term changes in the *Arabidopsis thaliana* glycerolipidome in response to temperature and light. *Plant J.* 66 (4), 656–668.
- Caird, M.A., Richards, J.H., Donovan, L.A., 2007. Nighttime stomatal conductance and transpiration in C<sub>3</sub> and C<sub>4</sub> plants. *Plant Physiol.* 143, 4–10.
- Campany, C.E., Tjoelker, M.G., von Caemmerer, S., Duursma, R.A., 2016. Coupled response of stomatal and mesophyll conductance to light enhances photosynthesis of shade leaves under sunflecks. *Plant Cell Environ.* 39, 2762–2773.
- Cardoso, A.A., Randall, J.M., McAdam, S.A., 2019. Hydraulics regulate stomatal responses to changes in leaf water status in the fern *Athyrium filix-femina*. *Plant Phys.* 179 (2), 533–543.
- Carins Murphy, M.R., Jordan, G.J., Brodribb, T.J., 2016. Cell expansion not cell differentiation predominantly co-ordinates veins and stomata within and among herbs and woody angiosperms grown under sun and shade. *Ann. Bot.* 118, 1127–1138.
- Carins Murphy, M.R., Dow, G.J., Jordan, G.J., Brodribb, T.J., 2017. Vein density is independent of epidermal cell size in *Arabidopsis* mutants. *Funct. Plant Biol.* 44 (4), 410–418.
- Carpita, N., Sabulase, D., Montezinos, D., Delmer, D.P., 1979. Determination of the pore size of cell walls of living plant cells. *Science* 205, 1144–1147.
- Carriqui, M., Roig-Oliver, M., Brodribb, T.J., Coopman, R., Gill, W., Mark, K., et al., 2019. Anatomical constraints to nonstomatal diffusion conductance and photosynthesis in lycophytes and bryophytes. *New Phytol.* 222, 1256–1270.
- Carriqui, M., Cabrera, H.M., Conesa, M.A., Coopman, R.E., Douthe, C., Gago, J., et al., 2015. Diffusional limitations explain the lower photosynthetic capacity of ferns as compared with angiosperms in a common garden study. *Plant Cell Environ.* 38, 448–460.
- Carriqui, M., Douthe, C., Molins, A., Flexas, J., 2019a. Leaf anatomy does not explain apparent short-term responses of mesophyll conductance to light and CO<sub>2</sub> in tobacco. *Physiol. Plant.* 165, 604–618.
- Casson, S.A., Hetherington, A.M., 2010. Environmental regulation of stomatal development. *Curr. Opin. Plant Biol.* 13, 90–95.
- Cathala, B., Monties, B., 2001. Influence of pectins on the solubility and the molar mass distribution of dehydrogenative polymers (DHPs, lignin model compounds). *Int. J. Biol. Macromol.* 29, 45–51.
- Chater, C.C.C., Caine, R.S., Fleming, A.J., Gray, J.E., 2017. Origins and evolution of stomatal development. *Plant Physiol.* 174, 624–638.
- Chaves, M.M., Flexas, J., Pinheiro, C., 2009. Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann. Bot.* 103, 551–560.
- Clemente-Moreno, M.J., Gago, J., Díaz-Vivancos, P., Bernal, A., Miedes, E., Bresta, P., Liakopoulos, G., Fernie, A.R., Hernández, J.A., Flexas, J., 2019. The apoplastic antioxidant system and altered cell wall dynamics influence mesophyll conductance and the rate of photosynthesis. *Plant J.* 99, 1031–1046.
- Cockburn, W., 1983. Stomatal mechanism as the basis of the evolution of CAM and C<sub>4</sub> photosynthesis. *Plant Cell Environ.* 6, 275–279.
- Cousins, A.B., Badger, M.R., von Caemmerer, S., 2006. A transgenic approach to understanding the influence of carbonic anhydrase on C1800 discrimination during C<sub>4</sub> photosynthesis. *Plant Physiol.* 142, 662–672.
- Daloso, D.M., Antunes, W.C., Pinheiro, D.P., Waquim, J.P., Araújo, W.L., Loureiro, M.E., Fernie, A.R., Williams, T.C.R., 2015. Tobacco guard cells fix CO<sub>2</sub> by both Rubisco and PEPcase while sucrose acts as a substrate during light-induced stomatal opening. *Plant Cell Environ.* 38, 2353–2371.
- Daloso, D.M., dos Anjos, L., Fernie, A.R., 2016a. Roles of sucrose in guard cell regulation. *New Phytol.* 211, 809–818.
- Daloso, D.M., Williams, T.C.R.R., Antunes, W.C., Pinheiro, D.P., Müller, C., Loureiro, M.E., Fernie, A.R., 2016b. Guard cell-specific upregulation of sucrose synthase 3 reveals that the role of sucrose in stomatal function is primarily energetic. *New Phytol.* 209, 1470–1483.
- Daloso, D.M., Medeiros, D.B., Dos Anjos, L., Yoshida, T., Araújo, W.L., Fernie, A.R., 2017. Metabolism within the specialized guard cells of plants. *New Phytol.* 216, 1018–1033.
- de Boer, H.J., Price, C.A., Wagner-Cremer, F., Dekker, S.C., Franks, P.J., Veneklaas, E.J., 2016a. Optimal allocation of leaf epidermal area for gas exchange. *New Phytol.* 210, 1219–1228.
- de Boer, H.J., Price, C.A., Wagner-Cremer, F., Dekker, S.C., Franks, P.J., Veneklaas, E.J., 2016b. Optimal allocation of leaf epidermal area for gas exchange. *New Phytol.* 210 (4), 1219–1228.
- de Souza Chaves, I., Feitosa-Araujo, E., Florian, A., Medeiros, D.B., da Fonseca-Pereira, P., Charton, L., et al., 2019d. The mitochondrial NAD<sup>+</sup> transporter (NDT1) plays important roles in cellular NAD<sup>+</sup> homeostasis in *Arabidopsis thaliana*. *Plant J.* 100, 487–504.
- Deans, R.M., Brodribb, T.J., Busch, F.A., Farquhar, G.D., 2019. Plant water-use strategy mediates stomatal effects on the light induction of photosynthesis. *New Phytol.* 222, 382–395.
- Desikan, R., Last, K., Harrett-Williams, R., Tagliavia, C., Harter, K., Hooley, R., et al., 2006. Ethylene-induced stomatal closure in *Arabidopsis* occurs via AtbHLH-mediated hydrogen peroxide synthesis. *Plant J.* 47, 907–916.
- Doheny-Adams, T., Hunt, L., Franks, P.J., Beerling, D.J., Gray, J.E., 2012. Genetic manipulation of stomatal density influences stomatal size, plant growth and tolerance to restricted water supply across a growth carbon dioxide gradient. *Philos. Trans. Biol. Sci.* 367, 547–555.
- Doi, M., Shimazaki, K., 2008. The stomata of the fern *Adiantum capillus-veneris* do not respond to CO<sub>2</sub> in the dark and open by photosynthesis in guard cells. *Plant Physiol.* 147, 922–930.
- Doi, M., Wada, M., Shimazaki, K.I., 2006. The fern *Adiantum capillus-veneris* lacks stomatal responses to blue light. *Plant Cell Physiol.* 47, 748–755.
- Douthe, C., Dreyer, E., Epron, D., Warren, C.R., 2011. Mesophyll conductance to CO<sub>2</sub>, assessed from online TDL-AS records of <sup>13</sup>CO<sub>2</sub> discrimination, displays small but significant short-term responses to CO<sub>2</sub> and irradiance in *Eucalyptus* seedlings. *J. Exp. Bot.* 62, 5335–5346.
- Douthe, C., Dreyer, E., Brendel, O., Warren, C.R., 2012. Is mesophyll conductance to CO<sub>2</sub> in leaves of three *Eucalyptus* species sensitive to short-term changes of irradiance under ambient as well as low O<sub>2</sub>? *Funct. Plant Biol.* 39, 435–448.
- Dow, G.J., Bergmann, D.C., 2014. Patterning and processes: how stomatal development defines physiological potential. *Curr. Opin. Plant Biol.* 21, 67–74.
- Duckett, J.G., Pressel, S., 2018. The evolution of the stomatal apparatus: intercellular spaces and sporophyte water relations in bryophytes—two ignored dimensions. *Philos. Trans. Biol. Sci.* 373.
- Edwards, D., Kerp, H., Hass, H., 1998. Stomata in early land plants: an anatomical and ecophysiological approach. *J. Exp. Bot.* 49, 255–278.
- Ellsworth, P.V., Ellsworth, P.Z., Koteyeva, N.K., Cousins, A.B., 2018. Cell wall properties in *Oryza sativa* influence mesophyll CO<sub>2</sub> conductance. *New Phytol.* 219, 66–76.
- Engelman, D.M., 2005. Membranes are more mosaic than fluid. *Nature* 438, 578.
- Engineer, C.B., Hashimoto-Sugimoto, M., Negi, J., Israelsson-Nordström, M., Azoulay-Shemer, T., Rappel, W.-J., et al., 2016a. CO<sub>2</sub> Sensing and CO<sub>2</sub> Regulation of Stomatal Conductance: Advances and Open Questions. *Trends Plant Sci.* 21, 16–30.
- Escudero, A., Del Arco, J.M., Garrido, M.V., 1992. The efficiency of nitrogen retranslocation from leaf biomass in *Quercus ilex* ecosystems. *Vegetatio* 99, 225–237.
- Evans, J.R., Sharkey, T.D., Berry, J.A., Farquhar, G.D., 1986. Carbon isotope discrimination measured concurrently with gas exchange to investigate CO<sub>2</sub> diffusion in leaves of higher plants. *Funct. Plant Biol.* 13, 281–292.
- Evans, J.R., Caemmerer, S.V., Satchell, B.A., Hudson, G.S., 1994. The relationship between CO<sub>2</sub> transfer conductance and leaf anatomy in transgenic tobacco with a reduced content of Rubisco. *Funct. Plant Biol.* 21, 475–495.
- Evans, J.R., Kaldenhoff, R., Genty, B., Terasima, I., 2009. Resistances along the CO<sub>2</sub> diffusion pathway inside leaves. *J. Exp. Bot.* 60, 2235–2248.
- Feild, T.S., Brodribb, T.J., 2013. Hydraulic tuning of vein cell microstructure in the evolution of angiosperm venation networks. *New Phytol.* 199 (3), 720–726.
- Finì, A., Loreto, F., Tattini, M., Giordano, C., Ferrini, F., Brunetti, C., Centritto, M., 2016. Mesophyll conductance plays a central role in leaf functioning of Oleaceae species exposed to contrasting sunlight irradiance. *Physiol. Plant.* 157, 54–68.
- Fleischer, A., O'Neill, M.A., Ehwald, R., 1999. The pore size of non-graminaceous plant cell walls is rapidly decreased by borate ester cross-linking of the pectic polysaccharide rhamnogalacturonan II. *Plant Physiol.* 121, 829–838.
- Flexas, J., Carriqui, M., 2020. Photosynthesis and photosynthetic efficiencies along the terrestrial plant's phylogeny: lessons for improving crop photosynthesis. *Plant J.* <https://doi.org/10.1111/tjp.14651>. (in press).
- Flexas, J., Díaz-Espejo, A., 2015. Interspecific differences in temperature response of mesophyll conductance: food for thought on its origin and regulation. *Plant Cell Environ.* 38, 625–628.



- Flexas, J., Bota, J., Escalona, J.M., Sampol, B., Medrano, H., 2002. Effects of drought on photosynthesis in grapevines under field conditions: an evaluation of stomatal and mesophyll limitations. *Funct. Plant Biol.* 29, 461–471.
- Flexas, J., Ribas-Carbó, M., Hanson, D.T., Bota, J., Otto, B., Cifre, J., McDowell, N., Medrano, H., Kaldenhoff, R., 2006. Tobacco aquaporin NtAQP1 is involved in mesophyll conductance to CO<sub>2</sub> *in vivo*. *Plant J.* 48, 427–439.
- Flexas, J., Diaz-Espejo, A., Galmés, J., Kaldenhoff, R., Medrano, H., Ribas-Carbó, M., 2007. Rapid variations of mesophyll conductance in response to changes in CO<sub>2</sub> concentration around leaves. *Plant Cell Environ.* 30, 1284–1298.
- Flexas, J., Barón, M., Bota, J., Ducruet, J.M., Gallé, A., Galmés, J., et al., 2009a. Photosynthesis limitations during water stress acclimation and recovery in the drought-adapted *Vitis* hybrid Richter-110 (*V. berlandieri* × *V. rupestris*). *J. Exp. Bot.* 60, 2361–2377.
- Flexas, J., Barón, M., Bota, J., Ducruet, J.M., Galle, A., Galmés, J., et al., 2009b. Photosynthesis limitations during water stress acclimation and recovery in the drought-adapted *Vitis* hybrid Richter-110 (*V. berlandieri* × *V. rupestris*). *J. Exp. Bot.* 60, 2361–2377.
- Flexas, J., Barbour, M.M., Brendel, O., Cabrera, H.M., Carriqui, M., Diaz-Espejo, A., et al., 2012. Mesophyll diffusion conductance to CO<sub>2</sub>: an unappreciated central player in photosynthesis. *Plant Sci.* 193, 70–84.
- Flexas, J., Gago, J., 2018. A role for ecophysiology in the 'omics' era. *Plant J.* 96, 251–259.
- Flexas, J., Niinemets, Ü., Gallé, A., Barbour, M.M., Centritto, M., Diaz-Espejo, A., et al., 2013a. Diffusional conductances to CO<sub>2</sub> as a target for increasing photosynthesis and photosynthetic water-use efficiency. *Phot. Res.* 117, 45–59.
- Flexas, J., Scoffoni, C., Gago, J., Sack, L., 2013b. Leaf mesophyll conductance and leaf hydraulic conductance: an introduction to their measurement and coordination. *J. Exp. Bot.* 64, 3965–3981.
- Flexas, J., Diaz-Espejo, A., Conesa, M.A., Coopman, R.E., Douthe, C., Gago, J., Gallé, A., Galmés, J., Medrano, H., Ribas-Carbó, M., Tomás, M., 2016. Mesophyll conductance to CO<sub>2</sub> and Rubisco as targets for improving intrinsic water use efficiency in C3 plants. *Plant Cell Environ.* 39, 965–982.
- Flexas, J., Carriqui, M., Cano, F.J., Coopman, R.E., Mizokami, Y., Tholen, D., Xiong, D., 2018a. CO<sub>2</sub> diffusion inside photosynthetic organs. in: WW Adams and I Terashima (eds), the leaf: a platform for performing photosynthesis. *Advances in Photosynthesis and Respiration* 44, 163–208.
- Flexas, J., Carriqui, M., Nadal, M., 2018b. Gas exchange and hydraulics during drought in crops: who drives whom? *J. Exp. Bot.* 69, 3791–3795.
- Flütsch, S., Wang, Y., Takemiya, A., Vialat-Chabrand, S., Klajchova, M., Nigro, A., Hills, A., Lawson, T., Blatt, M.R., Santelia, D., 2020. Guard Cell Starch Degradation Yields Glucose for Rapid Stomatal Opening in *Arabidopsis*. *Plant Cell*, Press.
- Foyer, C.H., 2018. Reactive oxygen species, oxidative signaling and the regulation of photosynthesis. *Environ. Exp. Bot.* 154, 134–142.
- Franco-Navarro, J.D., Rosales, M.A., Cubero-Font, P., Alvarez, R., Díaz-Espejo, A., Colmenero-Flores, J.M., 2019. Chloride as a macronutrient increases water-use efficiency by anatomically driven reduced stomatal conductance and increased mesophyll diffusion to CO<sub>2</sub>. *Plant J.* 99, 915–931.
- Franks, P.J., 2006. Higher rates of leaf gas exchange are associated with higher leaf hydrodynamic pressure gradients. *Plant Cell Environ.* 29, 584–592.
- Franks, P.J., Beerling, D.J., 2009. Maximum leaf conductance driven by CO<sub>2</sub> effects on stomatal size and density over geologic time. *PNAS* 106, 10343–10347.
- Franks, P.J., Britton-Harper, Z.J., 2016. No evidence of general CO<sub>2</sub> insensitivity in ferns: one stomatal control mechanism for all land plants? *New Phytol.* 211, 819–827.
- Franks, P.J., Farquhar, G.D., 2007. The mechanical diversity of stomata and its significance in gas-exchange control. *Plant Physiol.* 143, 78–87.
- Fromm, J., Fei, H., 1998. Electrical signalling and gas exchange in maize plants of drying soil. *Plant Sci.* 132, 203–213.
- Fujii, H., Chinnusamy, V., Rodrigues, A., Rubio, S., Antoni, R., Park, S.Y. et al. *In vitro* reconstitution of an abscisic acid signalling pathway. *Nature*. 462, 660–664.
- Fujita, T., Noguchi, K., Terashima, I., 2013. Apoplastic mesophyll signals induce rapid stomatal responses to CO<sub>2</sub> in *Commelina communis*. *New Phytol.* 199, 395–406.
- Fromm, J., Lautner, S., 2007. Electrical signals and their physiological significance in plants. *Plant Cell Environ.* 30, 249–257.
- Fujita, T., Noguchi, K., Ozaki, H., Terashima, I., 2019. Confirmation of mesophyll signals controlling stomatal responses by a newly devised transplanting method. *Funct. Plant Biol.* 46, 467–481.
- Gaastera, P., 1959. Photosynthesis of Crop Plants As Influenced by Light, Carbon Dioxide, Temperature, and Stomatal Diffusion Resistance (Doctoral Dissertation, Veenman).
- Gago, J., Douthe, C., Florez-Sarasa, I., Escalona, J.M., Galmés, J., Fernie, A.R., Flexas, J., Medrano, H., 2014. Opportunities for improving leaf water use efficiency under climate change conditions. *Plant Sci.* 226, 108–119.
- Gago, J., Daloso, D., de, M., Figueroa, C.M., Flexas, J., Fernie, A.R., Nikoloski, Z., 2016. Relationships of leaf net photosynthesis, stomatal conductance, and mesophyll conductance to primary metabolism: a multispecies meta-analysis approach. *Plant Physiol.* 171, 265–279.
- Gago, J., Carriqui, M., Nadal, M., Clemente-Moreno, M.J., Coopman, R.E., Fernie, A.R., Flexas, J., 2019. Photosynthesis optimized across land plant phylogeny. *Trends Plant Sci.* 24, 947–958.
- Gallé, A., Lautner, S., Flexas, J., Ribas-Carbó, M., Hanson, D., Roesgen, J., Fromm, J., 2013. Photosynthetic responses of soybean (*Glycine max* L.) to heat-induced electrical signalling are predominantly governed by modifications of mesophyll conductance for CO<sub>2</sub>. *Plant Cell Environ.* 36, 542–552.
- Gallé, A., Florez-Sarasa, I., El Aouad, H., Flexas, J., 2011. The Mediterranean evergreen *Quercus ilex* and the semi-deciduous *Cistus albidus* differ in their leaf gas exchange regulation and acclimation to repeated drought and re-watering cycles. *J. Exp. Bot.* 62 (14), 5207–5216.
- Gallé, A., Lautner, S., Flexas, J., Fromm, J., 2015. Environmental stimuli and physiological responses: the current view on electrical signalling. *Environ. Exp. Bot.* 114, 15–21.
- Galmés, J., Flexas, J., Savé, R., Medrano, H., 2007. Water relations and stomatal characteristics of Mediterranean plants with different growth forms and leaf habits: responses to water stress and recovery. *Plant Soil* 290, 139–155.
- Galmés, J., Conesa, M.A., Ochogavía, J.M., Perdomo, J.A., Francis, D.M., Ribas-Carbó, M., et al., 2011. Physiological and morphological adaptations in relation to water use efficiency in Mediterranean accessions of *Solanum lycopersicum*. *Plant Cell Environ.* 34, 245–260.
- Gan, Y., Zhou, L., Shen, Z.J., Shen, Z.X., Zhang, Y.Q., Wang, G.X., 2010. Stomatal clustering, a new marker for environmental perception and adaptation in terrestrial plants. *Bot. Stud.* 51.
- Gillon, J.S., Yakir, D., 2000. Naturally low carbonic anhydrase activity in C4 and C3 plants limits discrimination against C18O during photosynthesis. *Plant Cell Environ.* 23, 903–915.
- Gleason, S.M., Cooper, M., Wiggins, D.R., Bliss, C.A., Romay, M.C., Gore, M.A., et al., 2019. Stomatal conductance, xylem water transport, and root traits underpin improved performance under drought and well-watered conditions across a diverse panel of maize inbred lines. *Field Crops Res.* 234, 119–128.
- González-Guzmán, M., Pizzio, G.A., Antoni, R., Vera-Sirera, F., Merilo, E., Bassel, G.W., et al., 2012. Arabidopsis PYR/PYL/RCAR receptors play a major role in quantitative regulation of stomatal aperture and transcriptional response to abscisic acid. *Plant Cell* 24, 2483–2496.
- Gorton, H.L., Herbert, S.K., Vogelmann, T.C., 2003. Photoacoustic analysis indicates that chloroplast movement does not alter liquid-phase CO<sub>2</sub> diffusion in leaves of *Alocasia brisbanensis*. *Plant Physiol.* 132, 1529–1539.
- Gotow, K., Taylor, S., Zeiger, E., 1988. Photosynthetic carbon fixation in guard cell protoplasts of *Vicia faba* L. *Plant Physiol.* 86, 700–705.
- Grams, T.E.E., Koziolok, C., Lautner, S., Matussek, R., Fromm, J., 2007. Distinct roles of electric and hydraulic signals on the reaction of leaf gas exchange upon reirrigation in *Zea mays*. *Plant Cell Environ.* 30, 79–84.
- Granot, D., Kelly, G., 2019. Evolution of guard-cell theories: the story of sugars. *Trends Plant Sci.* 1–12.
- Gronidin, A., Rodrigues, O., Verdoucq, L., Merlot, S., Leonhardt, N., Maurel, C., 2015. Aquaporins contribute to ABA-triggered stomatal closure through OST1-mediated phosphorylation. *Plant Cell* 27, 1945–1954.
- Gu, L., Sun, Y., 2014a. Artefactual responses of mesophyll conductance to CO<sub>2</sub> and irradiance estimated with the variable J and online isotope discrimination methods. *Plant Cell Environ.* 37, 1231–1249.
- Gu, L., Sun, Y., 2014b. Artefactual responses of mesophyll conductance to CO<sub>2</sub> and irradiance estimated with the variable J and online isotope discrimination methods. *Plant Cell Environ.* 37, 1231–1249.
- Guanter, L., Zhang, Y., Jung, M., Joiner, J., Voigt, M., Berry, J., Frankenberg, C., Huete, A.R., Zarco-Tejada, P., Leeh, J., Morani, M.S., Ponce-Campos, G., Beer, C., Camps-Valls, G., Buchman, N., Gianelle, D., Klumpp, K., Cescattio, A., Baker, J.M., Griffis, T.J., 2014. Global and time-resolved monitoring of crop photosynthesis with chlorophyll fluorescence. *PNAS*. <https://doi.org/10.1073/pnas.1320081111>.
- Han, J., Lei, Z., Zhang, Y., Yi, X., Zhang, W., Zhang, Y., 2019. Drought-introduced variability of mesophyll conductance in *Gossypium* and its relationship with leaf anatomy. *Physiol. Plant.* 166, 873–887.
- Hanba, Y.T., Kogami, H., Terashima, I., 2002. The effect of growth irradiance on leaf anatomy and photosynthesis in *Acer* species differing in light demand. *Plant Cell Environ.* 25, 1021–1030.
- Hanba, Y.T., Shibasaki, M., Hayashi, Y., Hayakawa, T., Kasamo, K., Terashima, I., Katsuhara, M., 2004. Overexpression of the barley aquaporin HvPIP2; 1 increases internal CO<sub>2</sub> conductance and CO<sub>2</sub> assimilation in the leaves of transgenic rice plants. *Plant Cell Physiol.* 45, 521–529.
- Harley, P.C., Loreto, F., Di Marco, G., Sharkey, T.D., 1992. Theoretical considerations when estimating the mesophyll conductance to CO<sub>2</sub> flux by analysis of the response of photosynthesis to CO<sub>2</sub>. *Plant Physiol.* 98, 1429–1436.
- Harris, B.J., Harrison, C.J., Hetherington, A.M., Williams, T.A., 2020. Phylogenomic evidence for the monophyly of bryophytes and the reductive evolution of stomata. *Current Biology* 30 (11), 2001–2012.
- Harrison, E.L., Arce Cubas, L., Gray, J.E., Hepworth, C., 2019. The influence of stomatal morphology and distribution on photosynthetic gas exchange. *Plant J.* 101 (4), 768–779.
- Hashida, S.N., Itami, T., Takahashi, H., Takahara, K., Nagano, M., Kawai-Yamada, M., Shoji, K., Goto, F., Yoshihara, T., Uchimiya, H., 2010. Nicotinate/nicotinamide mononucleotide adenyltransferase-mediated regulation of NAD biosynthesis protects guard cells from reactive oxygen species in ABA-mediated stomatal movement in *Arabidopsis*. *J. Exp. Bot.* 61, 3813–3825.
- Haworth, M., Scutt, C.P., Douthe, C., Marino, G., Gomes, M.T.G., Loreto, F., et al., 2018. Allocation of the epidermis to stomata relates to stomatal physiological control: stomatal factors involved in the evolutionary diversification of the angiosperms and development of amphistomy. *Environ. Exp. Bot.* 151, 55–63.
- Heckwolf, M., Pater, D., Hanson, D.T., Kaldenhoff, R., 2011. The *Arabidopsis thaliana* aquaporin AtPIP1; 2 is a physiologically relevant CO<sub>2</sub> transport facilitator. *Plant J.* 67, 795–804.
- Hedrich, R., Geiger, D., 2017. Biology of SLAC1-type anion channels - from nutrient uptake to stomatal closure. *New Phytol.* 216, 46–61.
- Hedrich, R., Marten, I., 1993. Malate-induced feedback regulation of plasma membrane anion channels could provide a CO<sub>2</sub> sensor to guard cells. *EMBO J.* 12, 897–901.
- Henry, D.A., Simpson, R.J., Macmillan, R.H., 2000. Seasonal changes and the effect of temperature and leaf moisture content on intrinsic shear strength of leaves of pasture grasses. *Aust. J. Agric. Res.* 51, 823–831.



- Hetherington, A.M., Woodward, F.I., 2003. The role of stomata in sensing and driving environmental change. *Nature* 424, 901–908.
- Hite, D.R.C., Outlaw, W.H., Tarczynski, M.C., 1993. Elevated levels of both sucrose-phosphate synthase and sucrose synthase in *Vicia* guard cells indicate cell-specific carbohydrate interconversions. *Plant Physiol.* 101, 1217–1221.
- Ho, Q.T., Berghuijs, H.N., Watté, R., Verboven, P., Herremans, E., Yin, X., Retta, M.A., Aernouts, B., Saeys, W., Helfen, L., Farquhar, G.D., 2016. Three-dimensional micro-scale modelling of CO<sub>2</sub> transport and light propagation in tomato leaves enlightens photosynthesis. *Plant Cell Environ.* 39, 50–61.
- Hörak, H., Kollist, H., Merilo, E., 2017. Fern stomatal responses to ABA and CO<sub>2</sub> depend on species and growth conditions. *Plant Physiol.* 174, 672–679.
- Houston, K., Tucker, M.R., Chowdhury, J., Shirley, N., Little, A., 2016. The plant cell wall: a complex and dynamic structure as revealed by the responses of genes under stress conditions. *Fr. Plant Sci.* 7, 984.
- Hu, H., Boisson-Dernier, A., Israelsson-Nordström, M., Böhmer, M., Xue, S., Ries, A., et al., 2010. Carbonic anhydrases are upstream regulators of CO<sub>2</sub>-controlled stomatal movements in guard cells. *Nat. Cell Biol.* 12, 87–93.
- Hu, H., Rappel, W.J., Occhipinti, R., Ries, A., Böhmer, M., You, L., Xiao, C., Engineer, C.B., Boron, W.F., Schroeder, J.I., 2015. Distinct cellular locations of carbonic anhydrases mediate carbon dioxide control of stomatal movements. *Plant Physiol.* 169, 1168–1178.
- Hubbard, R.M., Ryan, M.G., Stiller, V., Sperry, J.S., 2001. Stomatal conductance and photosynthesis vary linearly with plant hydraulic conductance in ponderosa pine. *Plant Cell Environ.* 24, 113–121.
- Hughes, J., Hepworth, C., Dutton, C., Dunn, J.A., Hunt, L., Stephens, J., et al., 2017. Reducing stomatal density in barley improves drought tolerance without impacting on yield. *Plant Physiol.* 174, 776–787.
- Inoue, S., Kinoshita, T., 2017. Blue light regulation of stomatal opening and the plasma membrane H<sup>+</sup>-ATPase. *Plant Physiol.* 174, 531–538.
- Jang, J.Y., Kim, D.G., Kim, Y.O., Kim, J.S., Kang, H., 2004. An expression analysis of a gene family encoding plasma membrane aquaporins in response to abiotic stresses in *Arabidopsis thaliana*. *Plant Mol. Biol.* 54 (5), 713–725.
- Jarzyniak, K.M., Jasinski, M., 2014. Membrane transporters and drought resistance - a complex issue. *Fr. Plant Sci.* 5, 1–15.
- Jones, H.G., 1998. Stomatal control of photosynthesis and transpiration. *J. Exp. Bot.* 387–398.
- Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., et al., 2014. Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase BIK1 during plant immunity. *Mol. Cell* 54, 43–55.
- Kang, Y.U.N., Outlaw Jr., W.H., Andersen, P.C., Fiore, G.B., 2007. Guard-cell apoplastic sucrose concentration—a link between leaf photosynthesis and stomatal aperture size in the apoplastic phloem loader *Vicia faba* L. *Plant Cell Environ.* 30, 551–558.
- Kawase, M., Hanba, Y.T., Katsuhara, M., 2013. The photosynthetic response of tobacco plants overexpressing ice plant aquaporin McMIPB to a soil water deficit and high vapor pressure deficit. *J. Plant Res.* 126 (4), 517–527.
- Kelly, G., Moshelion, M., David-Schwartz, R., Halperin, O., Wallach, R., Attia, Z., et al., 2013. Hexokinase mediates stomatal closure. *Plant J.* 75, 977–988.
- Kenrick, P., Crane, P.R., 1997. The origin and early evolution of plants on land. *Nature* 389, 33.
- Khokh, M.A.R., Okuma, E.I.J.I., Hossain, M.A., Munemasa, S., Uraji, M., Nakamura, Y., et al., 2011. Involvement of extracellular oxidative burst in salicylic acid-induced stomatal closure in *Arabidopsis*. *Plant Cell Environ.* 34, 434–443.
- Kim, T.-H., Böhmer, M., Hu, H., Nishimura, N., Schroeder, J.I., 2010. Guard cell signal transduction network: advances in understanding abscisic acid, CO<sub>2</sub>, and Ca<sup>2+</sup> signalling. *Ann. Rev. Plant Biol.* 61, 561–591.
- Kjellbom, P., Larsson, C., 1984. Isolation of plasma membrane vesicles from leaves of spinach and barley, useful for studies on transport of carbon assimilation products. *Advances in Photosynthesis Research*. Springer, Dordrecht, pp. 673–676.
- Kjellbom, P., Larsson, C., Johansson, I., Karlsson, M., Johanson, U., 1999. Aquaporins and water homeostasis in plants. *Trends Plant Sci.* 4, 308–314.
- Knauer, J., Zaehle, S., De Kauwe, M.G., Bahar, N.H.A., Evans, J.R., Medlyn, B.E., Reichstein, M., Werner, C., 2019. Effects of mesophyll conductance on vegetation responses to elevated CO<sub>2</sub> concentrations in a land surface model. *Glob. Change Biol. Bioenergy* 25, 1820–1838.
- Knauer, J., Zaehle, S., De Kauwe, M.G., Haverd, V., Reichstein, M., Sun, Y., 2020. Mesophyll conductance in land surface models: effects on photosynthesis and transpiration. *Plant J.* 101 (4), 858–873.
- Kolbe, A.R., Brutnell, T.P., Cousins, A.B., Studer, A.J., 2018. Carbonic anhydrase mutants in *Zea mays* have altered stomatal responses to environmental signals. *Plant Physiol.* 177, 980–989.
- Kromdijk, J., Glowacka, K., Long, S.P., 2019. Photosynthetic efficiency and mesophyll conductance are unaffected in *Arabidopsis thaliana* aquaporin knock-out lines. *J. Exp. Bot.* 71, 318–329.
- Lake, J.A., Woodward, F.I., 2008. Response of stomatal numbers to CO<sub>2</sub> and humidity: control by transpiration rate and abscisic acid. *New Phytol.* 179, 397–404.
- Lampert, D.T.A., Kieliszewski, M.J., 2005. Stress upregulates periplasmic arabinogalactan-proteins. *Plant Biosystems-An International Journal Dealing with All Aspects of Plant Biology* 139, 60–64.
- Laporte, M.M., Shen, B., Tarczynski, M.C., 2002. Engineering for drought avoidance: expression of maize NADP-malic enzyme in tobacco results in altered stomatal function. *J. Exp. Bot.* 53, 699–705.
- Lawson, T., Blatt, M.R., 2014. Stomatal size, speed, and responsiveness impact on photosynthesis and water use efficiency. *Plant Physiol.* 164, 1556–1570.
- Lawson, T., Vialat-Chabrand, S., 2018. Speedy stomata, photosynthesis and plant water use efficiency. *New Phytol.* 221, 93–98.
- Lawson, T., Simkin, A.J., Kelly, G., Granot, D., 2014. Mesophyll photosynthesis and guard cell metabolism impacts on stomatal behaviour. *New Phytol.* 203, 1064–1081.
- Leakey, A.D., Lau, J.A., 2012. Evolutionary context for understanding and manipulating plant responses to past, present and future atmospheric CO<sub>2</sub>. *Philos. Trans. Biol. Sci.* 367, 613–629.
- Leakey, A.D., Ferguson, J.N., Pignon, C.P., Wu, A., Jin, Z., Hammer, G.L., Lobell, D.B., 2019. Water use efficiency as a constraint and target for improving the resilience and productivity of C3 and C4 crops. *Annu. Rev. Plant Biol.* 70, 781–808.
- Lee, A.G., 2004. How lipids affect the activities of integral membrane proteins. *Biochim. Biophys. Acta Biomembr.* 1666, 62–87.
- Li, S., Assmann, S.M., Albert, R., 2006. Predicting essential components of signal transduction networks: a dynamic model of guard cell abscisic acid signalling. *PLoS Biol.* 1732–1748.
- Li, Y., Xu, S., Gao, J., Pan, S., Wang, G., 2016. Glucose- and mannose-induced stomatal closure is mediated by ROS production, Ca<sup>2+</sup> and water channel in *Vicia faba*. *Physiol. Plant.* 252–261.
- Lima, V.F., Medeiros, D.B., Dos Anjos, L., Gago, J., Fennie, A.R., Daloso, D.M., 2018. Toward multifaceted roles of sucrose in the regulation of stomatal movement. *Plant Signal. Behav.* 00, 1–8.
- Lima, V.F., dos Anjos, L., Medeiros, D.B., Cândido-Sobrinho, S.A., Souza, L.P., Gago, J., et al., 2019. The sucrose-to-malate ratio correlates with the faster CO<sub>2</sub> and light stomatal responses of angiosperms compared to ferns. *New Phytol.* 223, 1873–1887.
- Loreto, F., Tsonev, T., Centritto, M., 2009. The impact of blue light on leaf mesophyll conductance. *J. Exp. Bot.* 60, 2283–2290.
- Lu, P., Zhang, S.Q., Outlaw, W.H., Riddle, K.A., 1995. Sucrose: a solute that accumulates in the guard-cell apoplast and guard-cell symplast of open stomata. *FEBS Lett.* 362, 180–184.
- Lu, P., Outlaw, Jr W.H., Smith, B.G., Freed, G.A., 1997. A new mechanism for the regulation of stomatal aperture size in intact leaves (accumulation of mesophyll-derived sucrose in the guard-cell wall of *Vicia faba*). *Plant Physiol.* 114, 109–118.
- Lu, Z., Lu, J., Pan, Y., Lu, P., Li, X., Cong, R., Ren, T., 2016. Anatomical variation of mesophyll conductance under potassium deficiency has a vital role in determining leaf photosynthesis. *Plant Cell Environ.* 39, 2428–2439.
- Ludwig, M., 2012. Carbonic anhydrase and the molecular evolution of C4 photosynthesis. *Plant Cell Environ.* 35, 22–37.
- Lugassi, N., Kelly, G., Fidel, L., Yaniv, Y., Attia, Z., Levi, A., et al., 2015. Expression of *Arabidopsis* hexokinase in *Citrus* guard cells controls stomatal aperture and reduces transpiration. *Fr. Plant Sci.* 6, 1–11.
- Males, J., Griffiths, H., 2017. Stomatal biology of CAM plants. *Plant Physiol.* 174, 550–560.
- Martins, S.C., Galmés, J., Cavatte, P.C., Pereira, L.F., Ventrella, M.C., DaMatta, F.M., 2014. Understanding the low photosynthetic rates of sun and shade coffee leaves: bridging the gap on the relative roles of hydraulic, diffusive and biochemical constraints to photosynthesis. *PLoS One* 9, e95571.
- Maurel, C., Verdoucq, L., Luu, D.T., Santoni, V., 2008. Plant aquaporins: membrane channels with multiple integrated functions. *Annu. Rev. Plant Biol.* 59, 595–624.
- McAdam, S.A.M., Brodribb, T.J., 2012a. Fern and lycophyte guard cells do not respond to endogenous abscisic acid. *Plant Cell* 24, 1510–1521.
- McAdam, S.A.M., Brodribb, T.J., 2012b. Stomatal innovation and the rise of seed plants. *Ecol. Lett.* 15, 1–8.
- McAdam, S.A.M., Brodribb, T.J., 2018. Mesophyll cells are the main site of abscisic acid biosynthesis in water-stressed leaves. *Plant Physiol.* 177, 911–917.
- McAdam, S.A.M., Brodribb, T.J., Banks, J.A., Hedrich, R., Atallah, N.M., Cai, C., et al., 2016. Abscisic acid controlled sex before transpiration in vascular plants. *PNAS* 113, 12862–12867.
- McAusland, L., Vialat-Chabrand, S., Davey, P., Baker, N.R., Brendel, O., Lawson, T., 2016. Effects of kinetics of light-induced stomatal responses on photosynthesis and water-use efficiency. *New Phytol.* 211, 1209–1220.
- McElwain, J.C., Yiotis, C., Lawson, T., 2016. Using modern plant trait relationships between observed and theoretical maximum stomatal conductance and vein density to examine patterns of plant macroevolution. *New Phytol.* 209, 94–103.
- Medeiros, D.B., Daloso, D.M., Fennie, A.R., Nikoloski, Z., Araújo, W.L., 2015. Utilizing systems biology to unravel stomatal function and the hierarchies underpinning its control. *Plant Cell Environ.* 38, 1457–1470.
- Medeiros, D.B., Martins, S.C.V., Cavalcanti, J.H.F., Daloso, D.M., Martinoia, E., Nunes-Nesi, A., et al., 2016. Enhanced photosynthesis and growth in *atac1* knockout mutants are due to altered organic acid accumulation and an increase in both stomatal and mesophyll conductance. *Plant Physiol.* 170, 86–101.
- Medeiros, D.B., Barros, K., Barros, J.A., Omena-Garcia, R.P., Arrivault, S., Pereira, Vincis, Sanglard, L., et al., 2017. Impaired malate and fumarate accumulation due the mutation of tonoplast dicarboxylate transporter. *Plant Physiol.* 175, 1068–1081.
- Medeiros, D.B., Perez-Souza, L., Antunes, W.C., Araújo, W.L., Daloso, D.M., Fennie, A.R., 2018. Sucrose breakdown within guard cells provides substrates for glycolysis and glutamine biosynthesis during light-induced stomatal opening. *Plant J.* 94, 583–594.
- Medeiros, D.B., da Luz, L.M., de Oliveira, H.O., Araújo, W.L., Daloso, D.M., Fennie, A.R., 2019. Metabolomics for understanding stomatal movements. *Theor. Exp. Plant Physiol.* 9, 91–102.
- Medlyn, B.E., Duursma, R.A., Eamus, D., Ellsworth, D.S., Prentice, I.C., Barton, C.V., et al., 2011. Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Glob. Change Biol. Bioenergy* 17, 2134–2144.
- Mercado, J.A., Matas, A.J., Heredia, A., Valpuesta, V., Quesada, M.A., 2004. Changes in the water binding characteristics of the cell walls from transgenic *Nicotiana tabacum* leaves with enhanced levels of peroxidase activity. *Physiol. Plant.* 122, 504–512.
- Merilo, E., Yarmolinsky, D., Jalakas, P., Parik, H., Tulva, I., Rasulov, B., Kollist, H., 2018. Stomatal VPD response: there is more to the story than ABA. *Plant Phys.* 176 (1), 851–864.
- Messinger, S.M., 2006. Evidence for involvement of photosynthetic processes in the

- stomatal response to CO<sub>2</sub>. *Plant Physiol.* 140, 771–778.
- Mittler, R., 2017. ROS are good. *Trends Plant Sci.* 22, 11–19.
- Miyazawa, S.I., Terashima, I., 2001. Slow chloroplast development in the evergreen broad-leaved tree species: relationship leaf anatomic characteristics and photosynthetic rate during leaf development. *Plant Cell Environ.* 24, 279–291.
- Miyazawa, S.I., Yoshimura, S., Shinzaki, Y., Maeshima, M., Miyake, C., 2008. Deactivation of aquaporins decreases internal conductance to CO<sub>2</sub> diffusion in tobacco leaves grown under long-term drought. *Funct. Plant Biol.* 35, 553–564.
- Mizokami, Y., Noguchi, K., Kojima, M., Sakakibara, H., Terashima, I., 2015. Mesophyll conductance decreases in the wild type but not in an ABA-deficient mutant (*aba1*) of *Nicotiana glauca* under drought conditions. *Plant Cell Environ.* 38, 388–398.
- Mizokami, Y., Noguchi, K., Kojima, M., Sakakibara, H., Terashima, I., 2019. Effects of instantaneous and growth CO<sub>2</sub> levels and abscisic acid on stomatal and mesophyll conductances. *Plant Cell Environ.* 42, 1257–1269.
- Momayyezi, M., Guy, R.D., 2017a. Substantial role for carbonic anhydrase in latitudinal variation in mesophyll conductance of *Populus trichocarpa* Torr. & Gray. *Plant Cell Environ.* 40, 138–149.
- Momayyezi, M., Guy, R.D., 2017b. Blue light differentially represses mesophyll conductance in high vs. Low latitude genotypes of *Populus trichocarpa* Torr. & Gray. *J. Plant Physiol.* 213, 122–128.
- Momayyezi, M., Guy, R.D., 2018. Concomitant effects of mercuric chloride on mesophyll conductance and carbonic anhydrase activity in *Populus trichocarpa* Torr. & Gray. *Trees* 32, 301–309.
- Momayyezi, M., McKown, A.D., Bell, S.C., Guy, R.D., 2020. Emerging roles for carbonic anhydrase in mesophyll conductance and photosynthesis. *Plant J.* 101, 831–844.
- Monties, B., 1989. Lignins. *Methods in Plant Biochemistry* Vol. 1. Academic Press, pp. 113–157.
- Moore, J.P., Nguema-Ona, E.E., Vreche-Gibouin, M., Sørensen, I., Willats, W.G., Driouch, A., Farrant, J.M., 2013. Arabinoxyle-rich polymers as an evolutionary strategy to plasticize resurrection plant cell walls against desiccation. *Planta* 237, 739–754.
- Mott, K.A., 2009. Opinion: stomatal responses to light and CO<sub>2</sub> depend on the mesophyll. *Plant Cell Environ.* 32, 1479–1486.
- Mott, K.A., Sibbersen, E.D., Shope, J.C., 2008. The role of the mesophyll in stomatal responses to light and CO<sub>2</sub>. *Plant Cell Environ.* 31, 1299–1306.
- Mott, K.A., Berg, D.G., Hunt, S.M., Peak, D., 2014. Is the signal from the mesophyll to the guard cells a vapour-phase ion? *Plant Cell Environ.* 37, 1184–1191.
- Nadal, M., Flexas, J., 2018. Mesophyll Conductance to CO<sub>2</sub>. *Water Scarcity and Sustainable Agriculture in Semiarid Environment: Tools, Strategies, and Challenges for Woody Crops*. pp. 403.
- Nadal, M., Flexas, J., 2019. Variation in photosynthetic characteristics with growth form in a water-limited scenario: implications for assimilation rates and water use efficiency in crops. *Agr. Wat. Manag.* 216, 457–472.
- Negi, J., Munemasa, S., Song, B., Tadakuma, R., Fujita, M., Azoulay-Shemer, T., et al., 2018. Eukaryotic lipid metabolic pathway is essential for functional chloroplasts and CO<sub>2</sub> and light responses in *Arabidopsis* guard cells. *PNAS* 115, 9038–9043.
- Niinemets, Ü., Díaz-Espejo, A., Flexas, J., Galmés, J., Warren, C.R., 2009. Role of mesophyll diffusion conductance in constraining potential photosynthetic productivity in the field. *J. Exp. Bot.* 60, 2249–2270.
- Nunes-Nesi, A., Carrari, F., Gibon, Y., Sulpice, R., Lytovchenko, A., Fisahn, J., et al., 2007. Deficiency of mitochondrial fumarate activity in tomato plants impairs photosynthesis via an effect on stomatal function. *Plant J.* 50, 1093–1106.
- Niinemets, Ü., Reichstein, Markus, 2003. Controls on the emission of plant volatiles through stomata: Differential sensitivity of emission rates to stomatal closure explained. *J. Geophys. Res.* 108 (D7). <https://doi.org/10.1029/2002JD002620>.
- Nunes-Nesi, A., Araújo, W.L., Fernie, A.R., 2011. Targeting mitochondrial metabolism and machinery as a means to enhance photosynthesis. *Plant Physiol.* 155, 101–107.
- O'Brien, J.A., Daudi, A., Butt, V.S., Bolwell, G.P., 2012. Reactive oxygen species and their role in plant defence and cell wall metabolism. *Planta* 236, 765–779.
- Onoda, Y., Wright, I.J., Evans, J.R., Hikosaka, K., Kitajima, K., Niinemets, Ü., et al., 2017. Physiological and structural tradeoffs underlying the leaf economics spectrum. *New Phytol.* 214, 1447–1463.
- Osborn, H.L., Alonso-Cantabrana, H., Sharwood, R.E., Covshoff, S., Evans, J.R., Furbank, R.T., von Caemmerer, S., 2017. Effects of reduced carbonic anhydrase activity on CO<sub>2</sub> assimilation rates in *Setaria viridis*: a transgenic analysis. *J. Exp. Bot.* 68, 299–310.
- Osborne, C.P., Beerling, D.J., Lomax, B.H., Chaloner, W.G., 2004. Biophysical constraints on the origin of leaves inferred from the fossil record. *PNAS* 101, 10360–10362.
- Otto, B., Uehlein, N., Siorra, S., Fischer, M., Ayaz, M., Belastegui-Macadam, X., Heckwolf, M., Lachnit, M., Pede, N., Priem, N., Reinhard, A., 2010. Aquaporin tetramer composition modifies the function of tobacco aquaporins. *J. Biol. Chem.* 285, 31253–31260.
- Outlaw, W.H., Manchester, J., Dicamelli, C.A., Randall, D.D., Rapp, B., Veith, G.M., 1979. Photosynthetic carbon reduction pathway is absent in chloroplasts of *Vicia faba* guard cells. *PNAS* 76, 6371–6375.
- Panchev, I.N., Slavov, A., Nikolova, Kr., Kovacheva, D., 2010. On the water-sorption properties of pectin. *Food Hydrocoll.* 24, 763–769.
- Park, S.Y., Fung, P., Nishimura, N., Jensen, D.R., Fujii, H., Zhao, Y., et al., 2009. Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* 324, 1068–1071.
- Peguero-Pina, J.J., Sisó, S., Flexas, J., Galmés, J., García-Nogales, A., Niinemets, Ü., Sancho-Knapik, D., Saz, M.A., Gil-Pelegrín, E., 2017a. Cell-level anatomical characteristics explain high mesophyll conductance and photosynthetic capacity in sclerophyllous Mediterranean oaks. *New Phytol.* 214, 585–596.
- Peguero-Pina, J.J., Sisó, S., Flexas, J., Galmés, J., Niinemets, Ü., Sancho-Knapik, D., Gil-Pelegrín, E., 2017b. Coordinated modifications in mesophyll conductance, photosynthetic potentials and leaf nitrogen contribute to explain the large variation in foliage net assimilation rates across *Quercus ilex* provenances. *Tree Physiol.* 37, 1084–1094.
- Penfield, S., Clements, S., Bailey, K.J., Gilday, A.D., Leegood, R.C., Gray, J.E., Graham, I.A., 2012. Expression and manipulation of PHOSPHOENOLPYRUVATE CARBOXYKINASE 1 identifies a role for malate metabolism in stomatal closure. *Plant J.* 69, 679–688.
- Perez-Martin, A., Michelazzo, C., Torres-Ruiz, J.M., Flexas, J., Fernandez, J.E., Sebastiani, L., Díaz-Espejo, A., 2014. Regulation of photosynthesis and stomatal and mesophyll conductance under water stress and recovery in olive trees: correlation with gene expression of carbonic anhydrase and aquaporins. *J. Exp. Bot.* 65, 3143–3156.
- Phillips, R., Ursell, T., Wiggins, P., Sens, P., 2009a. Emerging roles for lipids in shaping membrane-protein function. *Nature* 459, 379–385.
- Phillips, R., Ursell, T., Wiggins, P., Sens, P., 2009b. Emerging roles for lipids in shaping membrane-protein function. *Nature* 459, 379–385.
- Pickard, B.G., 1973. Action potentials in higher plants. *Bot. Rev.* 39, 172–201.
- Pieruschka, R., Huber, G., Berry, J.A., 2010. Control of transpiration by radiation. *PNAS* 107, 13372–13377.
- Pignocchi, C., Foyer, C.H., 2003. Apoplastic ascorbate metabolism and its role in the regulation of cell signalling. *Curr. Opin. Plant Biol.* 6, 379–389.
- Pitann, B., Schubert, S., Muhling, K.H., 2009. Decline in leaf growth under salt stress is due to an inhibition of H<sup>+</sup>-pumping activity and increase in apoplastic pH of maize leaves. *J. Plant Nutr. Soil Sci.* (1999) 172, 535–543.
- Pons, T.L., Flexas, J., Von Caemmerer, S., Evans, J.R., Genty, B., Ribas-Carbó, M., Brugnoli, E., 2009. Estimating mesophyll conductance to CO<sub>2</sub>: methodology, potential errors, and recommendations. *J. Exp. Bot.* 60, 2217–2234.
- Pou, A., Flexas, J., Alsina, M.D.M., Bota, J., Carambula, C., De Herralde, F., et al., 2008. Adjustments of water use efficiency by stomatal regulation during drought and recovery in the drought-adapted *Vitis* hybrid Richter-110 (*V. berlandieri* × *V. rupestris*). *Physiol. Plant.* 134, 313–323.
- Price, G.D., von Caemmerer, S., Evans, J.R., Yu, J.W., Lloyd, J., Oja, V., Kell, P., Harrison, K., Gallagher, A., Badger, M.R., 1994. Specific reduction of chloroplast carbonic anhydrase activity by antisense RNA in transgenic tobacco plants has a minor effect on photosynthetic CO<sub>2</sub> assimilation. *Planta* 193, 331–340.
- Qi, X., Torii, K.U., 2018. Hormonal and environmental signals guiding stomatal development. *BMC Biol.* 16, 1–11.
- Quirk, J., Andrews, M.Y., Leake, J.R., Banwart, S.A., Beerling, D.J., 2014. Ectomycorrhizal fungi and past high CO<sub>2</sub> atmospheres enhance mineral weathering through increased below-ground carbon-energy fluxes. *Biol. Lett.* 10, 20140375.
- Resco de Dios, V., Chowdhury, F.I., Granda, E., Yao, Y., Tissue, D.T., 2019. Assessing the potential functions of nocturnal stomatal conductance in C3 and C4 plants. *New Phytol.* 223, 1696–1706.
- Ritte, K., Rosenfeld, J., Rohrig, K., Raschke, K., 1999. Rates of sugar uptake by guard cell protoplasts of *Pisum sativum* L. Related to the solute requirement for stomatal opening. *Plant Physiol.* 121, 647–656.
- Robaina-Estévez, S., Daloso, D.M., Zhang, Y., Ferni, e A.R., Nikoloski, Z., 2017. Resolving the central metabolism of *Arabidopsis* guard cells. *Sci. Rep.* 7, 1–13.
- Rockwell, F.E., Holbrook, N.M., 2017. Leaf hydraulic architecture and stomatal conductance: a functional perspective. *Plant Phys.* 174 (4), 1996–2007.
- Roddy, A.B., Simonin, K.A., McCulloh, K.A., Brodersen, C.R., Dawson, T.E., 2018. Water relations of *Calycanthus* flowers: hydraulic conductance, capacitance, and embolism resistance. *Plant Cell Environ.* 41, 2250–2262.
- Rodrigues, O., Reshetnyak, G., Grondin, A., Saijo, Y., Leonhardt, N., Maurel, C., Verdoucq, L., 2017. Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA- and pathogen-triggered stomatal closure. *PNAS* 114, 9200–9205.
- Rodriguez-Dominguez, C.M., Buckley, T.N., Egea, G., de Cires, A., Hernandez-Santana, V., Martorell, S., Diaz-Espejo, A., 2016. Most stomatal closure in woody species under moderate drought can be explained by stomatal responses to leaf turgor. *Plant Cell Environ.* 39 (9), 2014–2026.
- Roig-Oliver, M., Nadal, M., Clemente-Moreno, M.J., Bota, J., Flexas, J., 2020. Cell wall components regulate photosynthesis and leaf water relations of *Vitis vinifera* cv. Grenache acclimated to contrasting environmental conditions. *J. Plant Physiol.* 244, 153084.
- Ros-Barceló, A., Gómez-Ros, L.V., Ferrer, M.A., Hernández, J.A., 2006. The apoplastic antioxidant enzymatic system in the wood-forming tissues of trees. *Trees-Struct. Funct.* 20, 145–156.
- Rusala, E.M., Beerling, D.J., Franks, P.J., Chater, C., Casson, S.A., Gray, J.E., Hetherington, A.M., 2011. Land plants acquired active stomatal control early in their evolutionary history. *Curr. Biol.* 21 (12), 1030–1035.
- Sack, L., Buckley, T.N., 2016. The developmental basis of stomatal density and flux. *Plant Physiol.* 171 (4), 2358–2363.
- Sack, L., Holbrook, N.M., 2006. Leaf hydraulics. *Annu. Rev. Plant Biol.* 57, 361–381.
- Sade, N., Shatil-Cohen, A., Attia, Z., Maurel, C., Boursiac, Y., Kelly, G., Granot, D., Yaaran, A., Lerner, S., Moshelion, M., 2014. The role of plasma membrane aquaporins in regulating the bundle sheath-mesophyll continuum and leaf hydraulics. *Plant Physiol.* 166, 1609–1620.
- Salles, S., Nardini, A., Pitt, F., Gullo, M.A.L., 2000. Xylem cavitation and hydraulic control of stomatal conductance in laurel (*Laurus nobilis* L.). *Plant Cell Environ.* 23, 71–79.
- Santelia, D., Lawson, T., 2016. Rethinking guard cell metabolism. *Plant Phys.* 172, 1371–1392.
- Santiago, J., Dupeux, F., Round, A., Antoni, R., Park, S.Y., Jamin, M., et al., 2009. The abscisic acid receptor PYR1 in complex with abscisic acid. *Nature* 462, 665–668.
- Sasaki, H., Hirose, T., Watanabe, Y., Ohsumi, R., 1998. Carbonic anhydrase activity and CO<sub>2</sub>-transfer resistance in Zn-deficient rice leaves. *Plant Phys.* 118, 929–934.
- Schiraldi, A., Fessas, D., Signorelli, M., 2012. Water activity in biological systems – a review. *Pol. J. Food Nutr. Sci.* 62, 5–13.
- Schmidt, R., Kunkowska, A.B., Schippers, J.H.M., 2016. Role of reactiveoxygen species

- during cell expansion in leaves. *Plant Physiol.* 172, 2098–2106.
- Scoffoni, C., Chatelet, D.S., Pasquet-kok, J., Rawls, M., Donoghue, M.J., Edwards, E.J., Sack, L., 2016. Hydraulic basis for the evolution of photosynthetic productivity. *Nat. Plants* 2, 1–8.
- Secchi, F., Perrone, I., Chitarra, W., Zwieniecka, A.K., Lovisolo, C., Zwieniecki, M.A., 2013. The dynamics of embolism refilling in abscisic acid (ABA)-deficient tomato plants. *Int. J. Mol. Sci.* 14, 359–377.
- Sharkey, T.D., Vassey, T.L., Vanderveer, P.J., Vierstra, R.D., 1991. Carbon metabolism enzymes and photosynthesis in transgenic tobacco (*Nicotiana tabacum* L.) having excess phytochrome. *Planta* 185, 287–296.
- Shatil-Cohen, A., Attia, Z., Moshelion, M., 2011. Bundle-sheath cell regulation of xylem-mesophyll water transport via aquaporins under drought stress: a target of xylem-borne ABA? *Plant J.* 67, 72–80.
- Simonin, K.A., Roddy, A.B., 2018. Genome downsizing, physiological novelty, and the global dominance of flowering plants. *PLoS Biol.* 16.
- Sobrado, M.A., 1992. The relationship between nitrogen and photosynthesis in relation to leaf age in a tropical xerophytic tree. *Photosynthetica* 26, 445–448.
- Sorrentino, G., Haworth, M., Wahbi, S., Mahmood, T., Zuomin, S., Centritto, M., 2016. Abscisic acid induces rapid reductions in mesophyll conductance to carbon dioxide. *PLoS One* 11 (2), e0148554.
- Studer, A.J., Gandin, A., Kolbe, A.R., Wang, L., Cousins, A.B., Brutnell, T.P., 2014. A limited role for carbonic anhydrase in C4 photosynthesis as revealed by a *calca2* double mutant in maize. *Plant Physiol.* 165, 608–617.
- Sun, Y., Gu, L., Dickinson, R.E., Norby, R.J., Pallardy, S.G., Hoffman, F.M., 2014. Impact of mesophyll diffusion on estimated global land CO<sub>2</sub> fertilization. *PNAS* 111 (44), 15774–15779.
- Sussmilch, F.C., Roelfsema, M.R.G., Hedrich, R., 2019a. On the origins of osmotically driven stomatal movements. *New Phytol.* 222, 84–90.
- Sussmilch, F.C., Schultz, J., Hedrich, R., Roelfsema, M.R.G., 2019b. Acquiring control: the evolution of stomatal signalling pathways. *Trends Plant Sci.* 24, 342–351.
- Syvtersen, J.P., Lloyd, J., McConchie, C., Kriedemann, P.E., Farquhar, G.D., 1995. On the relationship between leaf anatomy and CO<sub>2</sub> diffusion through the mesophyll of hypostomatous leaves. *Plant Cell Environ.* 18, 149–157.
- Takahashi, F., Suzuki, T., Osakabe, Y., Betsuyaku, S., Kondo, Y., Dohmae, N., et al., 2018. A small peptide modulates stomatal control via abscisic acid in long-distance signalling. *Nature* 556, 235.
- Talbott, L.D., Zeiger, E., 1993. Sugar and organic acid accumulation in guard cells of *Vicia faba* in response to red and blue light. *Plant Physiol.* 102, 1163–1.
- Tazoe, Y., von Caemmerer, S., Badger, M.R., Evans, J.R., 2009. Light and CO<sub>2</sub> do not affect the mesophyll conductance to CO<sub>2</sub> diffusion in wheat leaves. *J. Exp. Bot.* 60, 2291–2301.
- Tenhaken, R., 2015. Cell wall remodeling under abiotic stress. *Fr. Plant Sci.* 5, 771.
- Terashima, I., Ono, K., 2002. Effects of HgCl<sub>2</sub> on CO<sub>2</sub> dependence of leaf photosynthesis: evidence indicating involvement of aquaporins in CO<sub>2</sub> diffusion across the plasma membrane. *Plant Cell Physiol.* 43 (1), 70–78.
- Terashima, I., Hanba, Y.T., Tholen, D., Niinemets, Ü., 2011. Leaf functional anatomy in relation to photosynthesis. *Plant Physiol.* 155, 108–116.
- Thérout-Rancourt, G., Gilbert, M.E., 2017. The light response of mesophyll conductance is controlled by structure across leaf profiles. *Plant Cell Environ.* 40, 726–740.
- Thérout-Rancourt, G., Éthier, G., Pepin, S., 2015. Greater efficiency of water use in poplar clones having a delayed response of mesophyll conductance to drought. *Tree Physiol.* 35, 172–184.
- Tholen, D., Boom, C., Noguchi, K., Ueda, S., Katase, T., Terashima, I., 2008. The chloroplast avoidance response decreases internal conductance to CO<sub>2</sub> diffusion in *Arabidopsis thaliana* leaves. *Plant Cell Environ.* 31, 1688–1700.
- Tholen, D., Zhu, X.G., 2011. The mechanistic basis of internal conductance: a theoretical analysis of mesophyll cell photosynthesis and CO<sub>2</sub> diffusion. *Plant Physiol.* 156, 90–105.
- Tomás, M., Flexas, J., Copolovici, L., Galmés, J., Hallik, L., Medrano, H., et al., 2013. Importance of leaf anatomy in determining mesophyll diffusion conductance to CO<sub>2</sub> across species: quantitative limitations and scaling up by models. *J. Exp. Bot.* 64, 2269–2281.
- Tomás, M., Medrano, H., Brugnoli, E., Escalona, J.M., Martorell, S., Pou, A., et al., 2014. Variability of mesophyll conductance in grapevine cultivars under water stress conditions in relation to leaf anatomy and water use efficiency. *Austr. J. Grape Wine R* 20, 272–280.
- Tombesi, S., Nardini, A., Frioni, T., Soccolini, M., Zadra, C., Farinelli, D., et al., 2015. Stomatal closure is induced by hydraulic signals and maintained by ABA in drought-stressed grapevine. *Sci. Rep.* 5, 1–12.
- Tosens, T., Niinemets, Ü., Vislap, V., Eichelmann, H., Castro Diez, P., 2012a. Developmental changes in mesophyll diffusion conductance and photosynthetic capacity under different light and water availabilities in *Populus tremula*: how structure constrains function. *Plant Cell Environ.* 35, 839–856.
- Tosens, T., Niinemets, Ü., Westoby, M., Wright, I.J., 2012b. Anatomical basis of variation in mesophyll resistance in eastern Australian sclerophylls: news of a long and winding path. *J. Exp. Bot.* 63, 5105–5119.
- Tosens, T., Nishida, K., Gago, J., Coopman, R.E., Cabrera, H.M., Carriqui, M., et al., 2016. The Photosynthetic Capacity in 35 Ferns and Fern Allies: Mesophyll CO<sub>2</sub> Diffusion As a Key Trait.
- Uehlein, N., Lovisolo, C., Siefritz, F., Kaldenhoff, R., 2003. The tobacco aquaporin NtAQPI is a membrane CO<sub>2</sub> pore with physiological functions. *Nature* 425, 734–737.
- Uehlein, N., Otto, B., Hanson, D.T., Fischer, M., McDowell, N., Kaldenhoff, R., 2008. Function of *Nicotiana tabacum* aquaporins as chloroplast gas pores challenges the concept of membrane CO<sub>2</sub> permeability. *Plant Cell* 20 (3), 648–657.
- Uehlein, N., Kai, L., Kaldenhoff, R., 2017. Plant aquaporins and CO<sub>2</sub>. In: Chaumont, F., Tyerman, S. (Eds.), *Plant Aquaporins. From Transport to Signalling*. pp. 255–265.
- Vahisalu, T., Kollist, H., Wang, Y.F., Nishimura, N., Chan, W.Y., Valerio, G., et al., 2008. SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature* 452 (7186), 487–491.
- Verbančič, J., Lunn, J.E., Stitt, M., Persson, S., 2018. Carbon supply and the regulation of cell wall synthesis. *Mol. Plant* 11, 75–94.
- Veromann-Jürgenson, L.L., Tosens, T., Laanisto, L., Niinemets, Ü., 2017. Extremely thick cell walls and low mesophyll conductance: welcome to the world of ancient living! *J. Exp. Bot.* 68 (7), 1639–1653.
- Vrábl, D., Vaskova, M., Hronkova, M., Flexas, J., Santrucek, J., 2009. Mesophyll conductance to CO<sub>2</sub> transport estimated by two independent methods: effect of variable CO<sub>2</sub> concentration and abscisic acid. *J. Exp. Bot.* 60, 2315–2323.
- Vrábl, D., Vrábl, D., Hronková, M., Kubásek, J., Šantrůček, J., 2017. Stomatal function, density and pattern, and CO<sub>2</sub> assimilation in *Arabidopsis thaliana* *tmm1* and *sdd1-1* mutants. *Plant Biol.* 19, 689–701.
- Wang, S.W., Li, Y., Zhang, X.L., Yang, H.Q., Han, X.F., Liu, Z.H., et al., 2014a. Lacking chloroplasts in guard cells of crumpled leaf attenuates stomatal opening: both guard cell chloroplasts and mesophyll contribute to guard cell ATP levels. *Plant Cell Environ.* 37, 2201–2210.
- Wang, Y., Hills, A., Blatt, M.R., 2014b. Systems analysis of guard cell membrane transport for enhanced stomatal dynamics and water use efficiency. *Plant Physiol.* 164, 1593–1599.
- Wang, Y., Noguchi, K., Ono, N., Inoue, S., Terashima, I., Kinoshita, T., 2014c. Overexpression of Plasma Membrane H<sup>+</sup>-ATPase in Guard Cells Promotes Light-induced Stomatal Opening and.
- Wang, X., Du, T., Huang, J., Peng, S., Xiong, D., 2018. Leaf hydraulic vulnerability triggers the decline in stomatal and mesophyll conductance during drought in rice. *J. Exp. Bot.* 69, 4033–4045.
- Wang, H., Yan, S., Xin, H., Huang, W., Zhang, H., Teng, S., et al., 2019. A subsidiary cell-localized glucose transporter promotes stomatal conductance and photosynthesis. *Plant Cell* 31, 1328–1343.
- Watkins, J.M., Hechler, P.J., Muday, G.K., 2014. Ethylene-induced flavonol accumulation in guard cells suppresses reactive oxygen species and moderates stomatal aperture. *Plant Physiol.* 164, 1707–1717.
- Weraduwage, S.M., Campos, M.L., Yoshida, Y., Major, I.T., Kim, Y.S., Kim, S.J., Renna, L., Anozie, F.C., Brandizzi, F., Thomashow, M.F., Howe, G.A., 2018. Molecular mechanisms affecting cell wall properties and leaf architecture. *The Leaf: a Platform for Performing Photosynthesis*. Springer, Cham, pp. 209–253.
- Williams, T.G., Flanagan, L.B., Coleman, J.R., 1996. Photosynthetic gas exchange and discrimination against <sup>13</sup>CO<sub>2</sub> and C<sup>18</sup>O<sub>2</sub> in tobacco plants modified by an antisense construct to have low chloroplastic carbonic anhydrase. *Plant Physiol.* 112, 319–326.
- Willmer, C., Fricker, M., 1996. *Stomata*. Chapman & hall. London.
- Wooldenden, H.C., Baillie, A.L., Gray, J.E., Hobbs, J.K., Morris, R.J., Fleming, A.J., 2018. Models and mechanisms of stomatal mechanisms. *Trends Plant Sci.* 23 (9), 822–832.
- Wu, Y., Cosgrove, D.J., 2000. Adaptation of roots to low water potentials by changes in cell wall extensibility and cell wall proteins. *J. Exp. Bot.* 51, 1543–1553.
- WWAP (United Nations World Water Assessment Programme)/UN-Water, 2018. *The United Nations World Water Development Report 2018: Nature-based Solutions for Water*. UNESCO, Paris, pp. 22–23.
- Xiong, D., Douthe, C., Flexas, J., 2018. Differential coordination of stomatal conductance, mesophyll conductance, and leaf hydraulic conductance in response to changing light across species. *Plant Cell Environ.* 41, 436–450.
- Xiong, D., Liu, X., Liu, L., Douthe, C., Li, Y., Peng, S., et al., 2015. Rapid responses of mesophyll conductance to changes of CO<sub>2</sub> concentration, temperature and irradiance are affected by N supplements in rice. *Plant Cell Environ.* 38, 2541–2550.
- Xue, S., Hu, H., Ries, A., Merilo, E., Kollist, H., Schroeder, J.I., 2011. Central functions of bicarbonate in S-type anion channel activation and OST1 protein kinase in CO<sub>2</sub> signal transduction in guard cell. *EMBO J.* 30, 1645–1658.
- Yoshida, T., Anjos, L., dos Medeiros, D.B., Araújo, W.L., Fernie, A.R., Daloso, D.M., 2019a. Insights into ABA-mediated regulation of guard cell primary metabolism revealed by systems biology approaches. *Prog. Biophys. Mol. Bio.* 146, 37–49.
- Yoshida, T., Christmann, A., Yamaguchi-Shinozaki, K., Grill, E., Fernie, A.R., 2019b. Revisiting the basal role of ABA – roles outside of stress. *Trends Plant Sci.* 24, 625–635.
- Yoshida, T., Obata, T., Feil, R., Lunn, J.E., Fujita, Y., Yamaguchi-Shinozaki, K., Fernie, A.R., 2019c. The role of abscisic acid signalling in maintaining the metabolic balance required for *Arabidopsis* growth under nonstress conditions. *Plant Cell* 31, 84–105.
- Zhu, M., Zhu, N., Song, W.Y., Harmon, A.C., Assmann, S.M., Chen, S., 2014. Thiols-based redox proteins in abscisic acid and methyl jasmonate signalling in *Brassica napus* guard cells. *Plant J.* 78, 491–515. under elevated carbon dioxide. *Physiol. Plant.* 145, 395–405.
- Zoulas, N., Harrison, E.L., Casson, S.A., Gray, J.E., 2018. Molecular control of stomatal development. *Biochemical J.* 475 (2), 441–454.